



**UNIVERSIDADE FEDERAL DE SERGIPE**  
**PRÓ-REITORIA DE PÓS-GRADUAÇÃO E PESQUISA**  
**PROGRAMA DE PÓS-GRADUAÇÃO EM CIÊNCIAS DA SAÚDE**

**CRISTIANO DE QUEIROZ MENDONÇA**

**ALTERAÇÕES OFTALMOLÓGICAS EM PACIENTES PEDIÁTRICOS COM**  
**LEUCEMIA LINFOBLÁSTICA AGUDA**

**ARACAJU**

**2018**

**CRISTIANO DE QUEIROZ MENDONÇA**

**ALTERAÇÕES OFTALMOLÓGICAS EM PACIENTES PEDIÁTRICOS COM  
LEUCEMIA LINFOBLÁSTICA AGUDA**

Tese apresentada ao Programa de Pós-Graduação em  
Ciências da Saúde da Universidade Federal de  
Sergipe como requisito para obtenção de título de  
Doutor em Ciências da Saúde.

**Orientadora:** Profa Dra Rosana Cipolotti

**ARACAJU**

2018

FICHA CATALOGRÁFICA

FICHA CATALOGRÁFICA ELABORADA PELA BIBLIOTECA CENTRAL  
UNIVERSIDADE FEDERAL DE SERGIPE

M539a      Mendonça, Cristiano de Queiroz  
              Alterações oftalmológicas em pacientes pediátricos com  
              leucemia linfoblástica aguda / Cristiano de Queiroz Mendonça ;  
              orientadora Rosana Cipolotti. – Aracaju, 2018.  
              87 f. : il.

              Tese (doutorado em Ciências da Saúde) – Universidade  
              Federal de Sergipe, 2018.

              1. Leucemia linfoblástica. 2. Glaucoma. 3. Esteroides 4.  
              Quimioterapia. I. Cipolotti, Rosana, orient. II. Título.

CDU 616.155.392:616.7-053.2

**CRISTIANO DE QUEIROZ MENDONÇA**

**ALTERAÇÕES OFTALMOLÓGICAS EM PACIENTES PEDIÁTRICOS COM  
LEUCEMIA LINFOBLÁSTICA AGUDA**

Tese apresentada ao Programa de Pós-Graduação em  
Ciências da Saúde da Universidade Federal de Sergipe  
como requisito para obtenção de título de Doutor em  
Ciências da Saúde

Aprovado em 14/12/2018

**BANCA EXAMINADORA**

---

**Profa Dra. Rosana Cipolotti – Orientadora UFS**

---

**Prof. Dr. Marcelo Krieger Maestri – UFRGS**

---

**Prof. Dr. Kleyton de Andrade Bastos – UFS**

---

**Prof. Dr. Eduardo Luís de Aquino Neves– UFS**

---

**Prof. Dr. Bruno Campelo Leal – UNIT**

## RESUMO

**Introdução:** Leucemia Linfoblástica Aguda (LLA) é o câncer mais comumente encontrado entre os jovens, responsável por 26% dos casos de câncer infantil, com taxa de sobrevivência de doença de 90% em cinco anos. As manifestações oculares (MO) decorrentes das LLAs podem estar relacionadas à infiltração direta do olho e da órbita pelas células neoplásicas, serem secundárias às anormalidades vasculares tumor-induzidas ou a medicações usadas no tratamento como quimioterápicos e glicocorticoides (GC). Por ser doença oncológica com alto potencial de cura, em indivíduos jovens com elevada expectativa de vida, a identificação de eventuais complicações oculares de longo prazo decorrentes do tratamento e a correlação com fatores preditivos de recaída da doença poderá subsidiar o delineamento de um protocolo oftalmológico para esses casos. **Objetivo:** Caracterizar as MO em pacientes pediátricos com LLA e avaliar se estão associados a fatores de risco preditivos para recaída, com os protocolos (1999 ou 2009), gênero e infiltração líquórica. **Métodos:** Realizado estudo de coorte prospectivo em crianças e adolescentes com LLA, de janeiro de 2013 a dezembro de 2017, seguido de estudo de revisão sistemática, associando pacientes pediátricos em tratamento para LLA e hipertensão ocular (HO), devido a HO ser a MO mais prevalente. No estudo da coorte os pacientes foram submetidos a avaliações oftalmológicas antes do início do tratamento (D0), no oitavo dia (D8), no 28º dia (D28) e aos seis meses (D6 meses). A HO foi definida quando o resultado da aferição da pressão ocular foi  $>21$  mmHg. Medidas de acuidade visual (AV)  $<20/40$  foram consideradas como baixa visão (BAV). **Resultados:** Os resultados da coorte envolveram 55 pacientes e destes, 33% apresentaram MO. As principais foram HO (20%), hemorragia retiniana (7,3%) e BAV (7,3%). Forte associação foi encontrada entre pacientes com MO e alto risco de recaída ( $p = 0,035$ , Cramer V = 0,31) e os que usaram o protocolo de 1999 ( $p = 0,022$ , Cramer V = 0,32). O risco de MO em pacientes do protocolo de 1999 foi de 1,8 (IC = 1,154-2,804) e o risco de MO em pacientes de alto risco foi de 1,6 (IC = 1,111-2,442). Os resultados da revisão sistemática limitaram-se a quatro publicações sendo dois de relatos de casos individuais, um relato de cinco pacientes e outro descritivo prospectivo com doze pacientes, com resultados variando de total controle da pressão ocular e conservação da AV, até cegueira irreversível. **Conclusão:** Pacientes pediátricos com LLA têm alta incidência de MO devido ao tratamento e à própria doença, podendo ser assintomáticos ou evoluírem com complicações que podem chegar a BAV. Aqueles submetidos ao protocolo de 1999 e com alto risco de recaída são os mais propensos a apresentar MO e essas variáveis estão fortemente associadas. A HO é a MO mais prevalente. Poucos estudos foram encontrados correlacionando crianças com LLA e HO, com resultados variando de HO silenciosa, sem alterações visuais, até cegueira irreversível. Portanto, é proposto um protocolo que contemple exame oftalmológico sistemático com a medida da PIO imediatamente após o diagnóstico de LLA (D0) e, posteriormente, em D8, D28 e D6 meses.

**Descritores:** Leucemia linfoblástica aguda. Glaucoma. Esteroides. Quimioterapia.

## ABSTRACT

**Introduction:** Acute Lymphoblastic Leukemia (ALL) is a cancer commonly found among young people and responsible for 26% of childhood cancer cases. Today, up to 90% of such patients have an average survival rate of five-year free of the illness. Ocular manifestations (OM) arising from ALLs might be secondary to tumor-induced vascular anomalies or related to direct infiltration of the neoplastic cells into the eye and orbit, or to medications used in treatment such as chemotherapy and glucocorticoids (GC). Since this is an oncological disease with a high potential for cure, in young individuals with a high life expectancy, the identification of possible long-term ocular complications resulting from the treatment and correlation with predictive factors of relapse of the disease may support the design of an ophthalmological protocol for these patients cases. yet existent in scientific literature.

**Objective:** To characterize OM in pediatric patients under treatment for ALL and to evaluate if they are associated with predictive risk factors for relapse, with protocols (1999 or 2009), gender and cerebrospinal fluid (CSF) infiltration **Methods:** A prospective cohort study was conducted in children and adolescents with ALL from January 2013 to December 2017, followed by a systematic review associating pediatric patients under treatment for ALL and ocular hypertension (OH), because OH is the most prevalent OM. The patients underwent ophthalmologic evaluations before starting treatment (D0), on the eighth day (D8), at the 28th day (D28), and at six months (D6 months). Ocular hypertension (OH) was considered in results above 21 mmHg. Measures of visual acuity (VA) <20/40 were considered visual loss (VL). **Results:** The results of the cohort involved 55 patients and 18 (32.7%) presented OM, been OH (20.0%), retinal hemorrhage (7.3%) and VL (7.3%) the most frequent finds. A strong association was found between patients with OM and those with a high risk of relapse ( $p=0.035$ , Cramer  $V=0.31$ ) and who used the 1999 protocol ( $p=0.022$ , Cramer  $V=0.32$ ). The risk of OM in patients from the 1999 protocol was 1,799 (CI=1,154-2,804), while in those at high risk of relapse it was 1,647 (CI=1,111-2,442). The results of the systematic review were limited to a total of four publications, two of individual case reports, one report of five patients and another prospective descriptive with twelve patients, with results varying from total ocular pressure control and VA preservation, to irreversible blindness. **Conclusion:** Patients with ALL have a high incidence of OM due to the treatment and the disease itself, and it may even be asymptomatic and evolve with VL. Patients submitted to the 1999 protocol and at high risk of relapse are more likely to present OM and these variables are strongly associated. OH is the most prevalent OM. Few studies have been found correlating children with ALL and OH. Due to the possibility of disparate results from silent OH without visual changes to irreversible blindness. Therefore, a protocol is proposed that contemplates systematic ophthalmological examination with the measurement of IOP immediately after diagnosis of ALL (D0) and subsequently at D8, D28 and D6months.

**Descriptors:** acute lymphoblastic leukemia. Glaucoma. Steroids. Chemotherapy.

## LISTA DE ABREVIATURAS E SIGLAS

|                        |  |
|------------------------|--|
| <b>ALL</b>             | <i>Acute Lymphoblastic Leukemia</i>        |
| <b>AR</b>              | Alto risco de recaída da doença            |
| <b>BR</b>              | Baixo risco de recaída da doença           |
| <b>CSF</b>             | <i>Cerebrospinal fluid</i>                 |
| <b>D0</b>              | Dia anterior ao início do tratamento       |
| <b>D14</b>             | Décimo quarto dia de tratamento            |
| <b>D28</b>             | Vigésimo oitavo dia de tratamento          |
| <b>D8</b>              | Oitavo dia de tratamento                   |
| <b>D6meses</b>         | Sexto mês de tratamento                    |
| <b>GC</b>              | Glicocorticóides e <i>glucocorticoids</i>  |
| <b>GR</b>              | Gen receptor de glicocorticoide (GR-NR3C1) |
| <b>HO</b>              | Hipertensão Ocular                         |
| <b>IOP</b>             | <i>Intraocular pressure</i>                |
| <b>LLA</b>             | Leucemia Linfoblástica Aguda               |
| <b>LnH</b>             | Linfoma não-Hodgkin                        |
| <b>MO</b>              | Manifestações oculares                     |
| <b>MMP<sub>2</sub></b> | Matriz de metaloproteinase-2               |
| <b>NHL</b>             | <i>Non-Hodgkin Lymphoma</i>                |
| <b>OCT</b>             | Tomógrafo de Coerência Óptica              |
| <b>OD</b>              | Olho direito                               |
| <b>OE</b>              | Olho esquerdo                              |
| <b>OH</b>              | <i>Ocular hypertension</i>                 |
| <b>OM</b>              | <i>Ocular manifestations</i>               |
| <b>PIO</b>             | Pressão intraocular                        |
| <b>POAG</b>            | Glaucoma primário de ângulo aberto         |

|            |  |
|------------|--|
| <b>TM</b>  | Malha trabecular do ângulo da câmara anterior ocular |
| <b>TPA</b> | Activador do plasminogênio tecidual                  |
| <b>VA</b>  | <i>Visual acuity</i>                                 |
| <b>VL</b>  | <i>Visual loss</i>                                   |



## LISTA DE TABELAS

### ARTIGO 1

|                |  |    |
|----------------|--|----|
| <b>Table 1</b> | Profile of the patients included in the cohort.....  | 29 |
| <b>Table 2</b> | Comparison of patient groups with normal and altered ophthalmic examination for different variables..... | 30 |
| <b>Table 3</b> | Distribution of eye manifestations per patient.....  | 31 |
| <b>Table 4</b> | Relative risk of ocular manifestations.....  | 31 |

### ARTIGO 2

|                |   |    |
|----------------|---|----|
| <b>Table 1</b> | Design features and demographic details of studies included in the systematic review..... | 43 |
| <b>Table 2</b> | IOP control and complications in patients with steroid-induced ocular hypertension.....   | 44 |

## LISTA DE FIGURAS

### ARTIGO 1

- Figura 1** Flow chart showing the distribution of patients screened for possible ocular alterations in the cohort studied..... 28
- Figura 2** Boxplots showing the distribution of Intraocular Pressure in patients during the six-month follow-up period..... 32

### ARTIGO 2

- Figura 1** Flow diagram..... 43

## SUMÁRIO

|            |   |           |
|------------|---|-----------|
| <b>1</b>   | <b>INTRODUÇÃO.....</b>                                      | <b>10</b> |
| <b>2</b>   | <b>REVISÃO DA LITERATURA.....</b>                           | <b>11</b> |
| <b>2.1</b> | <b>MANIFESTAÇÕES OCULARES.....</b>                          | <b>13</b> |
| 2.1.1      | Manifestações Primárias.....                                | 13        |
| 2.1.2      | Manifestações Secundárias.....                              | 15        |
| 2.1.3      | Glaucoma Cortisônico e Hipertensão Ocular.....              | 16        |
| <b>3</b>   | <b>OBJETIVOS.....</b>                                       | <b>20</b> |
| <b>3.1</b> | <b>Objetivo Geral.....</b>                                  | <b>20</b> |
| <b>3.2</b> | <b>OBJETIVOS ESPECÍFICOS.....</b>                           | <b>20</b> |
| <b>4</b>   | <b>CASUÍSTICA E MÉTODOS.....</b>                            | <b>21</b> |
| <b>4.1</b> | <b>CASUÍSTICA.....</b>                                      | <b>21</b> |
| 4.1.1      | Modelo de Estudo, População, Amostra e Local do estudo..... | 21        |
| 4.1.2      | Critérios de Inclusão para o Estudo.....                    | 21        |
| 4.1.3      | Critérios de Exclusão para o Estudo.....                    | 21        |
| <b>4.2</b> | <b>MÉTODOS.....</b>   | <b>22</b> |
| 4.2.1      | Estudo de Coorte (Artigo 1) .....                           | 22        |
| 4.2.2      | Estudo de Revisão Sistemática (Artigo 2) .....              | 23        |
| 4.2.3      | Considerações Éticas.....                                   | 23        |
| <b>5</b>   | <b>RESULTADOS.....</b>                                      | <b>24</b> |
| <b>5.1</b> | <b>Estudo de Coorte (Artigo 1) .....</b>                    | <b>24</b> |
| <b>5.2</b> | <b>Estudo de Revisão Sistemática (Artigo 2) .....</b>       | <b>37</b> |
| <b>5.3</b> | <b>Regras para Publicação.....</b>                          | <b>45</b> |
| 5.3.1      | Estudode Coorte (Artigo 1) .....                            | 45        |
| 5.3.2      | Estudo de Revisão Sistemática (Artigo 2) .....              | 59        |
| <b>6</b>   | <b>DISCUSSÃO GERAL.....</b>                                 | <b>67</b> |
| <b>7</b>   | <b>CONCLUSÃO.....</b>                                       | <b>72</b> |
| <b>7.1</b> | <b>Conclusão dos Artigos.....</b>                           | <b>72</b> |
| 7.1.1      | ARTIGO 1 - Estudo de Coorte.....                            | 72        |
| 7.1.2      | ARTIGO 2 -Estudo de Revisão Sistemática.....                | 73        |

|                                 |           |
|---------------------------------|-----------|
| <b>REFERÊNCIAS.....</b>         | <b>74</b> |
| <b>PRODUÇÃO CIENTÍFICA.....</b> | <b>81</b> |
| <b>ANEXO A.....</b>             | <b>83</b> |
| <b>APÊNDICE A.....</b>          | <b>84</b> |



## 1 INTRODUÇÃO

A leucemia linfoblástica aguda (LLA) é, na faixa pediátrica, o câncer mais comum e responde por aproximadamente 26% dos casos de câncer infantil (WARD et al., 2014). A LLA corresponde a 80% dos casos diagnosticados em pacientes de zero a 14 anos e aproximadamente 56% dos casos em adolescentes acima dos 14 anos. Nos últimos 30 anos, os avanços científicos se traduziram em melhorias nas propostas terapêuticas que culminaram numa redução da mortalidade por LLA em mais de 30% (WARD et al., 2014). Atualmente tais pacientes possuem, em média, sobrevida livre de doença de até 90% em cinco anos (MARCOUX et al., 2016; WARD et al., 2014).

Apesar do sucesso do tratamento quimioterápico para LLA, vários estudos apontam para alterações tardias em diversos órgãos e sistemas, incluindo os domínios emocional e cognitivo (ALLEMANI et al., 2015; BERBIS et al., 2014; HONG et al., 2014; MERTENS et al., 2014; WHELAN et al., 2010). Lesões oculares e orbitais são a terceira manifestação extramedular mais comum da LLA, depois das lesões meníngeas e testiculares, variando entre 43 a 90% dos casos de LLA (CHARIF et al., 2002, SHARMA et al., 2004; TALCOTT; GARG, R.; GARG, S., 2016).

As MO tanto podem estar relacionadas à infiltração direta do olho e da órbita pelas células neoplásicas, quanto serem secundárias à anormalidades vasculares tumor-induzida ou a complicações pelo uso de medicamentos quimioterápicos, incluindo altas doses de glicocorticoides (GC) (MENDONCA et al., 2014; RAZEGHINEJAD; KATZ, 2012).

Por se tratar de uma doença oncológica com grande potencial de cura e que compromete indivíduos com elevada expectativa de vida, é importante caracterizar as MO de pacientes pediátricos em tratamento para LLA, bem como avaliar se tais manifestações possuem correlação com preditores de risco de recidiva definidos na literatura e outros fatores como protocolo de tratamento quimioterápico, idade, gênero, óbito, imunofenotipagem e infiltração do liquor por células neoplásicas. Identificar e tratar precocemente tais alterações oculares pode prevenir dano permanente à visão, além de diagnosticar precocemente uma possível infiltração ou recidiva incipiente da doença.

## 2 REVISÃO DA LITERATURA

A LLA constitui 26% de todas as neoplasias em indivíduos de zero a 15 anos, constituindo 80% das leucemias infantis. Seu pico de incidência ocorre entre dois e cinco anos de idade, com média de idade de 6,5 anos, sendo discretamente mais frequente no sexo masculino, em indivíduos de cor branca e de países desenvolvidos (WARD et al., 2014; YEOH et al., 2013; YOUNG et al., 1986). História familiar está presente em 5% dos casos (YASMEEN; ASHRAF, 2009; YOUNG et al., 1986).

As taxas de LLA variaram cerca de 10 vezes entre 54 populações estudadas, sendo mais elevadas em hispânicos brancos dos EUA, seguido pelo Equador, e sendo mais baixas na população negra dos EUA, Tunísia e Uganda. Essa variação quanto à distribuição geográfica pode ser atribuída a fatores de risco ambientais, genética e/ou diferenças nos critérios de diagnóstico e no tratamento (LINET et al., 2015). Estudos epidemiológicos de câncer na América do Sul não são comumente relatados. A mediana da taxa de incidência ajustada de leucemia no Brasil e na Argentina foi de 47 por milhão (DE CAMARGO et al., 2010; MORENO et al., 2013). No Brasil, em Recife-PE, estudo estimou em cerca de 48,5 casos por milhão de crianças e adolescentes (LINS et al., 2016).

Em virtude dos avanços no tratamento, a taxa de mortalidade da LLA declinou mais de 30% nos últimos 30 anos (WARD et al., 2014). Por se tratar de uma neoplasia maligna das células precursoras de linfócitos, caracterizada pela aberração na proliferação e diferenciação dos precursores linfoides, a LLA acarreta falência do sistema imunológico e decréscimo na hematopoese normal, resultando em anemia, trombocitopenia e neutropenia. É comum ao exame a presença de anemia (86%), linfadenopatia (75%), hepatomegalia (67%) e esplenomegalia (58%). Inicialmente pode haver leucocitose elevada ( $>50.000/\text{mm}^3$ ), observada em 34% dos pacientes, hemoglobina  $<7\text{g/dl}$  em 54%, plaquetopenia  $<20.000/\text{mm}^3$  em 33% e comprometimento de sistema nervoso central em 5% (YASMEEN; ASHRAF, 2009).

Os pacientes são classificados no momento do diagnóstico em dois grupos, de acordo com o risco de recidiva. Os pacientes com alto risco de recaída (AR) são aqueles que preencheram um dos seguintes critérios: idade igual ou superior a nove anos, contagem de leucócitos maior que ou igual a  $50.000\text{ células}/\text{mm}^3$  no momento do diagnóstico e

imunofenotipagem compatível com LLA de células T. O grupo de baixo risco (BR) engloba os pacientes que não preenchem nenhum dos critérios acima (CERDÀ-IBÁÑEZ et al., 2018).

Existem diferenças notáveis entre os países de baixa e média renda em comparação com os países desenvolvidos em relação aos índices de mortalidade durante a indução, mortalidade aos seis meses após o diagnóstico (início da fase de manutenção), infiltração no líquido cefalorraquidiano e proporção de pacientes alto risco de recaída. Essas diferenças estão mais diretamente relacionadas ao acesso à infraestrutura adequada para o tratamento de complicações do que aos aspectos biológicos da doença (HOWARD et al., 2008).

Hoje, nos país desenvolvidos, uma criança com diagnóstico de LLA tem uma expectativa de sobrevida média em cinco anos de até 90%, com 95% de probabilidade de viver mais 10 anos passados os cinco iniciais (MARCOUX et al., 2016; WARD et al., 2014). A Sociedade Americana de Câncer mostra que em 2016, 6590 novos casos foram diagnosticados, com 1400 mortes secundárias à LLA (TERWILLIGER; ABDUL-HAY, 2017).

O tratamento consiste em quatro a seis semanas de quimioterapia de indução inicialmente administrada no hospital, seguida por vários meses de quimioterapia de consolidação (geralmente até os seis meses do início do tratamento) e dois a três anos de quimioterapia de manutenção (MARGOLIN et al., 2010). Os mais recentes protocolos de tratamento da LLA utilizados no Brasil foram propostos em 1999 e atualizados em 2009 (ALL-99 e ALL-09) pelo “Grupo Brasileiro de Tratamento da Leucemia Linfoblástica Aguda na Infância” da Sociedade Brasileira de Oncologia Pediátrica. Ao final dos primeiros 28 dias de tratamento (D28) o protocolo ALL-99 previa uma dosagem total de 1120 mg/m<sup>2</sup> de Prednisona para o grupo BR e 1680 mg/m<sup>2</sup> de Prednisona para o grupo AR. O protocolo ALL-09 previa 420 mg/m<sup>2</sup> de prednisona e 126 mg/m<sup>2</sup> de dexametasona para o grupo BR e 1260 mg/m<sup>2</sup> de prednisona para o grupo AR (BRANDALISE et al., 2010; WATANABE, 2007).

Apesar do sucesso do tratamento quimioterápico para LLA, vários estudos apontam para alterações tardias em diversos órgãos e sistemas, incluindo os domínios emocional e cognitivo, deficiência de crescimento, risco aumentado para o aparecimento de novo câncer como Leucemia Mieloide Aguda ou Linfoma, tumor de sistema nervoso central e de pescoço (BERBIS et al., 2014; HONG et al., 2014; MERTENS et al., 2014; WARD et al., 2014; WHELAN et al., 2010). Lesões oculares e orbitais são a terceira manifestação extramedular mais comum da leucemia aguda, depois das lesões meníngeas e testiculares (CHARIF et al., 2002; GORDON et al., 2001).



## 2.1 Manifestações Oculares

De maneira geral, a fisiopatologia do acometimento ocular na LLA deve-se a três mecanismos principais: I) infiltração direta do olho e órbita pelas células neoplásicas; II) anormalidades vasculares afetando a retina ou III) acometimento neuro-oftalmológico (REDDY; MENON, 1998). No primeiro caso pode-se ter invasão de coroide, hifema, hipópio, heterocromia da íris, glaucoma secundário, proptose, episclerite (ÇAÇA et al., 2005; YALCINBAYIR et al., 2017); no segundo, hemorragias retinianas, manchas de Roth, exsudatos algodinosos, oclusões vasculares, descolamento de retina, microaneurismas, dilatação e tortuosidade venosa (ÇAÇA et al., 2005; CHAUDHURI; ROY, S.; ROY, P., 2013; YALCINBAYIR et al., 2017); no terceiro, a infiltração do nervo óptico pode acarretar paralisia de nervos cranianos e papiledema (SHARMA et al., 2004). Entre as anormalidades retinianas, uma revisão observou 24% de hemorragias retinianas, 11% de manchas de Roth e 16% de exsudatos (TALCOTT; GARG, R.; GARG, S., 2016). Mais frequente parece ser a infiltração da coroide, estimada em 50 a 82% dos casos (GORDON et al., 2001).

Envolvimento ocular pode ser classificado em duas principais categorias: primária ou infiltração leucêmica direta, e secundária ou envolvimento indireto (SHARMA et al., 2004). A infiltração leucêmica direta pode ser observada em três diferentes padrões: (a) infiltração uveal, infiltração orbital, e sinais neuro-oftalmológicos de infiltração nervosa (CHAUDHURI; ROY, S.; ROY, P., 2013; LIN et al., 2005), (b) paralisia de nervos cranianos e (c) papiledema (NGUYEN; HAIDER; ACKERMAN, 2013). As alterações secundárias são manifestações retinianas, hemorragia vítrea, infecções, e oclusões vasculares devido as anormalidades hematológicas da leucemia como anemia, trombocitopenia, hiperviscosidade e imunodepressão (SHUYUAN; MING; YUNXIA, 2018).

### 2.1.1 Manifestações primárias

A córnea é uma estrutura avascular e, portanto, não é comumente envolvida na leucemia, especialmente na forma de invasão direta de células leucêmicas. Pode haver envolvimento da córnea além do envolvimento do limbo (ALLEN; STRAATSMA, 1961). Alterações corneanas são também vistas quando o seu epitélio se transforma devido ao

tratamento quimioterápico. Essas mudanças incluem adelgaçamento irregular, maturação defeituosa e queratinização (JABS et al., 1983). Apesar de o envolvimento da conjuntiva não ser uma apresentação comum da leucemia, ocorre mais comumente em pacientes em leucemias linfóides (DUKE-ELDER, 1965).

Infiltração da esclera é geralmente um achado de autópsia e ocorre em leucemias agudas. Essas células são achadas mais frequentemente na episclera em um padrão perivascular (ALLEN; STRAATSMA, 1961).

Clinicamente, infiltração evidente da íris por células leucêmicas não é comum. Isso ocorre com o envolvimento da coroide e do corpo ciliar e é clinicamente caracterizado por mudança na cor da íris e um pseudohipópio, de coloração cinza/amarelada (PERRY; MALLIN, 1979). Leucemias foram identificadas como a doença de base em 5% dos casos de uveítes na pediatria (SOYLU; OZDEMIR; ANLI, 1997).

A coroide demonstra infiltração leucêmica mais consistentemente no exame histopatológico, embora, clinicamente, a retina seja a parte mais envolvida na leucemia. O envolvimento da coroide por células leucêmicas tende a ser perivascular e desigual ou difuso (ALLEN; STRAATSMA, 1961).

O envolvimento retiniano é estimado em até 69% de todos os pacientes com leucemia mostram alteração de fundo de olho em algum ponto durante o curso da doença (ALEMAYEHU et al., 1996). As manifestações precoces, por conta dos distúrbios hematológicos, são dilatação e tortuosidade venosa (BALLANTYNE; MICHAELSON, 1970). As hemorragias e infiltrações são vistas em todos os níveis da retina, especialmente nas camadas interiores com destruição focal. Os infiltrados e agregados de células leucêmicas são geralmente vistos em torno das áreas hemorrágicas. A membrana limitante interna frequentemente atua como uma eficiente barreira contra a infiltração de células leucêmicas. (KUWUBARA; AIELLO, 1964). Contudo, células leucêmicas ocasionalmente invadem o vítreo possivelmente emergindo da cabeça do nervo óptico (REESE; GUY, 1976).

Como consequência do aumento na sobrevida, o envolvimento do sistema nervoso e do nervo óptico se tornou mais frequente, particularmente na leucemia aguda. Pode aparecer, mesmo quando a medula óssea está em remissão, já que a barreira hematoencefálica restringe a livre passagem de certos agentes quimioterápicos (RIDGWAY; JAFFE; WALTON, 1976; SHAW et al., 1960). Infiltração do sistema nervoso central ocorre tanto em adulto, menos comum, quanto em crianças (DAWSON; ROSENTHAL; MOLONEY, 1973) e mais comumente em LLA comparado às Leucemias Mielóides Agudas (HYMAN et al., 1965).

Infiltração orbital na leucemia se apresenta como exoftalmia, edema palpebral e dor (CAVDAR et al., 1971; COLOMBINI et al., 1995). Todos os tipos de leucemia podem atingir a órbita, porém o envolvimento orbital é mais comum nas leucemias agudas comparadas as crônicas e ocorrem mais comumente em leucemias linfóides quando comparadas as mielóides (JAKOBIEC; JONES, 1979).

Outras alterações oculares menos comuns da leucemia incluem necrose do segmento anterior, dacriocistite e infiltração da pele (CULLIS; HINES; BULLOCK, 1952).

### 2.1.2 Manifestações Secundárias

Pacientes com leucemia, durante o período de neutropenia, são mais suscetíveis a desenvolver infecções incomuns e que ameacem a vida. Esses pacientes são suscetíveis a uma ampla variedade de infecções virais, fúngicas, bacterianas e de protozoários (COGAN, 1977). Uma das infecções mais comuns nos imunocomprometidos é por citomegalovírus (CMV) (SHIBATA et al., 1997), que invade a retina, causando necrose, hemorragia, e combina descolamento exsudativo e regmatogênico da retina (MEREDITH; AABERG; REESER, 1979). Outros vírus (herpes simples, varicela zoster e caxumba) também podem causar retinite necrotizante em pacientes imunocomprometidos (COGAN, 1977). Herpes zoster também pode causar úlcera corneal periférica, ceratite e esclerite (WALTON; REED, 1999).

Os fungos são a causa mais comum de infecções oculares nas leucemias. Infecção por *Candida* causa uveíte e retinite com infiltratos algodinosos no vítreo. Outra infecção fúngica comum em pacientes com leucemia é por *Aspergillus* (COGAN, 1977).

Infecção bacteriana por *Pseudomonas* é comum em pacientes imunossuprimidos. Essa infecção pode iniciar como blefarconjuntivite e se estender causando celulite orbitária (GIAGOUNIDIS et al., 1997).

Sintomas oculares podem desenvolver-se após transplante da medula óssea, tanto pela doença enxerto *versus* hospedeiro como pela radioterapia de corpo inteiro, que integra protocolos de condicionamento que antecedem o transplante de medula óssea. Melhorias recentes no manejo geral desses pacientes têm resultado no reconhecimento mais frequente dos problemas oculares. MO da síndrome enxerto *versus* hospedeiro incluem ceratoconjuntivite sicca, lagoftalmo cicatricial, conjuntivite por *pseudomonas* ou estéril, defeitos epiteliais, úlcera corneana, uveíte e ectrópio da pálpebra (CLAES; KESTELYN, 2000; KASMANN; RUPRECHT, 1993).

Nos últimos anos foram introduzidas muitas novas drogas anticancerosas, tanto na prática clínica quanto como parte de protocolos de pesquisa. Para muitos destes, a ampla gama de efeitos colaterais ainda não é conhecida. Bussulfan, por exemplo, foi identificado como causa de catarata subcapsular posterior (RAVINDRANATHAN; PAUL; KURIAKOSE, 1972). Vincristina e vinblastina são tóxicos ao sistema nervoso e afetam principalmente os nervos periféricos, podendo acometer os motores ocular e do trigêmeo. Vincristina também foi apontada como causa de atrofia óptica (SHURIN; REKATE; ANNABLE, 1982).

Os GC, por sua vez, usados em dose imunossupressora durante a fase de indução e manutenção, aumentam significativamente o risco de o paciente desenvolver catarata subcapsular posterior, seu principal efeito colateral. Segundo estudo, esse risco é cinco vezes maior quando comparados os pacientes curados com seus irmãos saudáveis (ESSIG et al., 2014). Outro efeito importante é a Hipertensão Ocular (HO) e glaucoma iatrogênico pelo uso de GC, o glaucoma cortisônico (KERSEY; BROADWAY, 2005; MENDONÇA et al., 2014; RAZEGHINEJAD; KATZ, 2012).

### 2.1.3 Glaucoma Cortisônico e Hipertensão Ocular

A descoberta dos GC foi um grande avanço no tratamento de várias doenças. À semelhança de outros agentes terapêuticos, têm seus próprios efeitos colaterais, incluindo HO e glaucoma iatrogênico (RAZEGHINEJAD; KATZ, 2012).

O glaucoma é a principal causa de cegueira irreversível, mas prevenível, do mundo. Tem como principal causa o aumento da Pressão Intra Ocular (PIO) (POMORSKA et al., 2012), que é mais frequentemente aferida pela técnica de tonometria de aplanção após instilação de anestesia local (proparacaina a 0,5%) e corante de fluoresceína (GOLDMAN, 1957).

A incidência de glaucoma na infância é de 2,29/100 000 pessoas com idade inferior a 20 anos. O glaucoma infantil é uma patologia pediátrica incomum, associada a significativa perda visual. Consiste de um grupo heterogêneo de doenças que geram neuropatia óptica e perda de campo visual, e que pode ser classificado em primário, secundário e adquirido (APONTE; DIEHL; MOHNEY, 2010).

Glaucoma primário é geralmente dividido em congênito (cujo início se dá desde o nascimento até o final do período de recém-nascido) e primário juvenil de ângulo aberto (cujo início se dá mais frequentemente dos quatro anos até o período de adulto jovem). O glaucoma

primário congênito é o principal tipo observado na infância (APONTE; DIEHL; MOHNEY, 2010). Ainda no grupo dos glaucomas congênitos, os secundários associam-se a alterações sindrômicas ou a outras condições presentes ao nascimento, como aniridia, síndrome de Axenfeld-Rieger, retinopatia da prematuridade, síndrome de Rubinstein-Taybi, síndrome de Sturge-Weber, persistência do vítreo hiperplástico primário e rubéola congênita (APONTE; DIEHL; MOHNEY, 2010). Glaucoma adquirido é resultado de outros processos não presentes ao nascimento como inflamação, drogas (como GC), trauma e cirurgia (APONTE; DIEHL; MOHNEY, 2010).

HO induzida por GC foi primeiramente descrita em 1950 com a observação de glaucoma em associação com a administração de GC sistêmico (MCLEAN, 1950). As rotas mais comuns de indução HO são a administração tópica, intra-ocular ou periocular. Pode também ocorrer depois de uso sistêmico, aplicação na pele, intranasal, ou por inalação (URBAN; DREYER, 1993). Estudo prévio mostrou possibilidade de HO em valores elevados, em criança com Síndrome Nefrótica, usuária de GC (BRITO et al., 2012). Geralmente ocorre com algumas semanas de uso de GC em paciente susceptíveis, mas é geralmente reversível com a descontinuação do tratamento. Entretanto, se o tratamento for prolongado, pode resultar em neuropatia óptica (NG et al, 2008; SPAETH; BARROS; FUDEMBERG, 2009; QASHTON, 2011). As alterações glaucomatológicas comprometem não somente o nervo óptico como também interferem nos mecanismos que causam elevação da PIO, como aumentando a resistência à drenagem do humor aquoso pela malha trabecular (TM) (CLARK; WORDINGER, 2009; DANIAS et al., 2011). Além disso, o aumento da PIO pode ser pela própria infiltração da leucemia na malha trabecular (ROWAN; SLOAM, 1976).

Assim, o uso de GC podem acarretar HO e gerar uma patologia semelhante ao Glaucoma Crônico de Ângulo Aberto (POAG) (SCHWARTZ et al., 2002). Estudos em pacientes tratados com potentes GC tópicos ou sistêmicos mostraram o desenvolvimento de HO em 30 a 40% dos casos, pela diminuição da drenagem do humor aquoso associada a modificações morfológicas na TM (ARMALY; BECKER, 1965) devido ao maior acúmulo de proteínas como a fibronectina e miocilina além da degradação do ativador do plasminogênio tecidual (TPA), e/ou matriz de metaloproteinase-2 (MMP<sub>2</sub>) responsáveis pela homeostasia do sistema de drenagem do humor aquoso na câmara anterior, TM e consequentemente aumento da PIO (STAMER et al., 2013). Em adição, foi observado que pacientes mais sensíveis ao uso de GC possuem susceptibilidade maior para desenvolver POAG. Esses pacientes apresentam nível elevado de cortisol sérico e metabolismo anormal de cortisol ocular (CLARK; WORDINGER, 1995).

A administração de GC pode aumentar a PIO em 90% dos pacientes com POAG. Avaliando-se as pessoas normotensas oculares na população geral, a probabilidade de aumento da PIO com uso de GC é de 30-40%, caracterizando os médios-responsivos. PIO acima de 31 mm Hg ou 15 mm Hg acima do valor de base ocorre entre 4-6% dos casos, para os altos-responsivos (ZHANG; CLARK; YORIO, 2005). Até 5% dos pacientes não respondem ao tratamento medicamentoso e necessitam cirurgia (RAZEGHINEJAD; KATZ, 2012).

Embora o glaucoma seja também observado em síndrome de Cushing, desencadeada pela produção de excesso endógeno de GC, a probabilidade de elevação da PIO pela administração de GC por via sistêmica é menor do que por via tópica. Geralmente os grupos de pacientes que estão sendo tratados com GC sistêmicos em longo prazo apresentaram maiores médias e maiores picos de valores de PIO que a população normal (HOVLAND; ELLIS, 1967; LEE, 1958). Um estudo encontrou uma incidência de 10% de glaucoma em pacientes transplantados renais que receberam GC (ADHIKARY; SELLS; BASU, 1982). Há correlação positiva entre a PIO e dose de GC (aumento de 1,4 mmHg na PIO média para cada aumento de 10 mg na dose diária média de prednisona administrada) (TRIPATHI et al., 1992).

Vários relatos de casos clínicos mostram aumento da PIO após o uso de GC em crianças (BRITO et al., 2012; THAM et al., 2004; YAMASHITA et al., 2010). Com a suspensão do tratamento a pressão normalmente retorna a níveis normais. Por outro lado, o uso de GC sistêmicos pode gerar aumento assintomático da PIO, inclusive entre os pacientes pediátricos (NG et al., 2008).

Há sugestões da existência de diferenças genéticas entre os pacientes responsivos e não-responsivos. Alguns autores tentam explicar a susceptibilidade genética por mecanismo monozigótico autossômico: heterozigóticos para os médios-responsivos e os homozigóticos para os altos-responsivos (RAZEGHINEJAD; KATZ, 2011). Diversos mecanismos diferentes podem ser responsáveis para as sensibilidades GC entre os indivíduos, incluindo polimorfismos no gene GR (NR3C1), gene regulador do GC, e as diferenças na expressão do GR. A resposta ao GC é regulada pelos níveis relativos do receptor alfa (GR $\alpha$ ) e do regulador negativo (GR $\beta$ ). Linhagens de células da TM de pacientes glaucomatosos têm uma relação GR $\beta$ -GR $\alpha$  menor em comparação com as células TM normais, tornando-os mais sensíveis aos GCs (STAMER et al., 2013). Em situações de homeostase há produção normal de miocilina, fibronectina, glicosaminas e laminas, e degradação na TM do ativador do plasminogênio tecidual (TPA), e/ou matriz de metaloproteinase-2 (MMP<sub>2</sub>) mediados pelo

GR $\beta$ -GR $\alpha$ . Quando há aumento da relação GR $\alpha$ -GR $\beta$  ocorre produção aumentada de miocilina e conseqüentemente menor drenagem de humor aquoso, predispondo ao aumento da PIO (PFEFFER et al., 2010; STAMER et al., 2013). A diminuição de GR $\beta$  pode resultar no aumento da resposta ao GC e aumento da PIO (STAMER et al., 2013).

Estudos prévios têm mostrado que alterações morfológicas no nervo óptico e nas camadas de fibras nervosas do nervo óptico podem preceder alterações de campo de visão em pacientes com glaucoma (POMORSKA et al., 2012).

O conhecimento sobre as manifestações oculares da leucemia é importante para o diagnóstico e tratamento da doença, além de muitas vezes refletir o estágio da mesma no organismo. Estudos demonstram que a evidência de envolvimento ocular pode estar associada a pior prognóstico (CURTO et al., 1989; KINCAID; GREEN, 1983; REDDY; MENON, 1998).

Poucos estudos foram publicados especificamente relacionando MO em pacientes pediátricos tratados para LLA e nenhum deles foi elaborado de maneira sistemática e prospectiva. A identificação de eventuais complicações oculares de longo prazo decorrentes da própria doença e do tratamento, além de avaliar se as MO possuem correlação com preditores de risco de recidiva da doença e outros fatores (protocolo de tratamento quimioterápico, idade, gênero, óbito, imunofenotipagem e infiltração do CSF por células neoplásicas), poderá subsidiar o delineamento de um protocolo oftalmológico para esses casos.

### **3 OBJETIVOS**

#### **3.1 Objetivo Geral**

- Avaliar as MO em pacientes pediátricos portadores de LLA.

#### **3.2 Objetivos Específicos**

- Identificar possíveis MO em pacientes pediátricos com LLA (artigo 1).
- Correlacionar se fatores prognósticos estabelecidos para recidiva de doença oncológica (tipo de protocolo de tratamento quimioterápico utilizado, sexo e infiltração do líquido por células neoplásicas) podem prever risco de desenvolver MO (artigo 1).
- Descrever a produção científica nacional e internacional em forma de Revisão Sistemática da MO mais prevalente em pacientes pediátricos com LLA, a HO (artigo 2).



## **4 CASUÍSTICA E MÉTODOS**

### **4.1 Casuística**

#### **4.1.1 Modelo de estudo, população, amostra e local do estudo**

Foi desenvolvido um estudo de coorte prospectiva, realizado em pacientes pediátricos com diagnóstico de LLA, tratadas no período de 1 de julho de 2013 a 30 de dezembro de 2017 no Centro de Oncologia Dr. Oswaldo Leite (serviço público especializado em oncologia pediátrica do estado de Sergipe, região Nordeste do Brasil). Os pacientes elegíveis foram incluídos consecutivamente, após confirmação citopatológica e imunofenotípica do diagnóstico e após a anuência do responsável para o estudo de campo.

Realizou-se um estudo de Revisão Sistemática associando a MO mais prevalente em crianças com LLA, a HO.

#### **4.1.2 Critérios de Inclusão para o Estudo**

- a) Diagnóstico de LLA confirmado laboratorialmente.

- b) Idade até 19 anos incompletos.
- c) Ausência de tratamento quimioterápico anterior.
- d) Ausência de outra patologia por ocasião do diagnóstico.
- e) Não utilização de GC sistêmico nos seis meses que antecederam o diagnóstico.

#### 4.1.3 Critérios de Exclusão para o Estudo

- a) Impossibilidade técnica.
- b) Diagnóstico prévio compatível com glaucoma ou doença anterior relacionada a qualquer mudança na PIO.

## 4.2 Métodos

### 4.2.1 Estudo de Coorte (Artigo 1)

Pacientes elegíveis foram classificados ao diagnóstico em dois grupos conforme o risco de recaída da doença: pacientes com alto risco de recaída (AR): idade igual ou maior que nove anos, contagem de leucócitos maior ou igual a 50.000 células/mm<sup>3</sup> no momento do diagnóstico e imunofenotipagem compatível com LLA de células T. O grupo de baixo risco de recaída (BR) compreendeu aqueles não inseridos em nenhum desses critérios.

Os protocolos de tratamento de LLA utilizados foram os propostos pelo “Grupo Brasileiro para tratamento da Leucemia Linfóide Aguda na Infância” em 1999 e atualizado em 2009 (ALL-99 e ALL-09). Foi utilizado como rotina de tratamento, o protocolo ALL-99 em todos pacientes admitidos até 31 de dezembro de 2014 e o ALL-09 após essa data.

Todos os pacientes foram submetidos a exame oftalmológico realizados por um único oftalmologista (C.Q.M.) conforme as técnicas e a periodicidade descritas a seguir. As avaliações foram efetuadas nos dias zero (D0), oitavo (D8), vigésimo oitavo (D28) dia do início do tratamento e com seis meses de tratamento (D6meses).

Os exames de D0 e D8 foram realizados no leito hospitalar devido à fragilidade da saúde desses pacientes quando do diagnóstico e consistiram em avaliação de segmento anterior ocular com lâmpada de fenda portátil, segmento posterior com oftalmoscopia direta

sem midríase e medida da PIO. Os exames de (D28) e (D6meses) foram realizados em um consultório oftalmológico especializado, sendo a análise de segmento anterior por biomicroscopia com lâmpada de fenda, o de segmento posterior através de oftalmoscopia indireta sob midríase induzida por tropicamida 0,01%, aferida a Acuidade Visual (AV) e a medida da PIO.

A PIO foi aferida por tonometria de aplanção com aparelho tipo Perkins, após anestesia local com colírio de proparacaína a 0,5% e corante de fluoresceína (GOLDMANN, 1957). Foram consideradas compatíveis com HO as medidas de PIO acima de 21 mmHg (CURTO et al., 1989).

A AV foi aferida com projetor automatizado de optotipos, com e sem midríase. Foi utilizado a tabela de Snellen e considerado baixa visual quando  $AV < 20/40$ .

Os pacientes diagnosticados com qualquer patologia ocular receberam tratamento específico apropriado.

A análise dos dados foi feita utilizando-se o programa SPSS (*Statistical Package for Social Sciences*) 24.0. Variáveis categóricas foram descritas como frequência absolutas e relativas comparadas usando o teste exato de Fischer com o teste Cramer V para força e teste de ajuste multivariável de Benjamini-Hochberg para falsos positivos. As variáveis contínuas foram descritas em termos de média e desvio padrão. Foi utilizado também o teste Binomial para comparar proporções entre grupos não-paramétricos e o teste de Pearson para correlação de variáveis com distribuição normal. Foram considerados estatisticamente significativos os valores de  $p < 0,05$ .

#### 4.2.2 Estudo de Revisão Sistemática (Artigo 2)

Um protocolo da revisão sistemática foi registrado previamente no banco de dados PROSPERO (número de registro CRD42018087146).

Pesquisa bibliográfica foi realizada nas bases de dados: PubMed, Scopus, Web of Science, Science Direct, e Cochrane Library até 31 de janeiro de 2018. Uma pesquisa na literatura cinzenta foi realizada através do Google Scholar e OpenThesis. Os primeiros 100 resultados do Google Scholar foram registrados. A pesquisa também incluiu uma busca manual de referências cruzadas de artigos e resenhas originais.

A exposição foi definida como tratamento de LLA ou LNH com GC sistêmicos, independentemente do tipo de GC ou dosagem. O desfecho foi a HO ocorrida após o início do GC e definida como uma PIO superior a 21 mmHg em um ou ambos os olhos.

#### 4.2.3 Considerações Éticas

O projeto foi aprovado pelo Comitê de Ética em Pesquisa envolvendo Seres Humanos da Universidade Federal de Sergipe (CEP-UFS), com número: 214.759. (APÊNDICE A)

## 5 RESULTADOS

### 5.1 Estudo de Coorte (Artigo 1)

#### **Ocular Manifestations in Acute Lymphoblastic Leukemia: A Five-Year Cohort Study of Pediatric Patients**

(Artigo aceito para publicação no Jornal: Leukemia Research, Manuscript Number: LR-D-18-00551R1).

#### **ABSTRACT**

**Objective:** To characterize ocular manifestations (OM) of pediatric patients treating for acute lymphoblastic leukemia (ALL) and to evaluate whether they are associated with well-described predictive risk factors for relapse, protocol (1999 or 2009), gender and cerebrospinal fluid infiltration. **Methods:** A prospective cohort study was conducted in children and adolescents with ALL from January 2013 to December 2017. The patients underwent ophthalmologic evaluations before starting treatment (D0), on the eighth day (D8), at the 28th day (D28), and at six months (D6 months). Ocular hypertension (OH) was

considered in results above 21 mmHg. Measures of visual acuity <20/40 were considered visual loss (VL) **Results:** Fifty-five patients were examined and 18 (32.7%) presented OM, been OH (61.1%), retinal hemorrhage (22.2%) and VL (22.2%) the most frequent finds. A strong association was found between patients with OM and those with a high risk of relapse ( $p=0.035$ , Cramer  $V=0.31$ ) and who used the 1999 protocol ( $p=0.022$ , Cramer  $V=0.32$ ). The risk of OM in patients from the 1999 protocol was 1,799 (CI=1,154-2,804), while in those at high risk of relapse it was 1,647 (CI=1,111-2,442). **Conclusions:** Patients with ALL have a high incidence of OM due to the treatment and the disease itself, and it may even be asymptomatic and evolve with VL. Of these, we can highlight OH as the most prevalent. Patients submitted to the 1999 protocol and at high risk of relapse are more likely to present OM and these variables are strongly associated.

**Key words:** acute lymphoblastic leukemia, glaucoma, steroids, chemotherapy

## INTRODUCTION

Ocular manifestations in Acute Lymphoblastic Leukemia (ALL), varying from 43% to 90% depending on the study, are concerning because they are often silent [1,2]. Such manifestations often go unperceived since most patients are asymptomatic. Nevertheless, they can indicate a relapse or early worsening of the condition with a potential risk to the patient's sight.

Generally speaking, the physiopathology of ocular impairment in ALL is attributed to three main mechanisms: I) direct infiltration of the eye and orbit by the neoplastic cells; II) vascular abnormalities affecting the retina or III) neuro-ophthalmic impairment [3]. In the first case, there could be an invasion of the choroid, hyphemia, hypopyon, heterochromia of the iris, secondary glaucoma, proptosis and episcleritis [4,5]; in the second case, retinal hemorrhages, Roth spots, cotton wool exudation, vascular occlusions, retinal detachment, microaneurysms, as well as venal dilatation and tortuosity [4,5,6]; in the third case, infiltration of the optic nerve can lead to paralysis of cranial nerves and papilledema [2]. One revision observed retinal hemorrhages at 24%, Roth spots at 11% and exudates at 16% [1]. The most frequent appears to be an infiltration of the choroid, ranging from 50% to 82% of the cases [7].

Furthermore, the effects of chemotherapy on the patients' vision cannot be ignored. The use of high doses of glucocorticosteroids (GC) in the treatment entails a significantly

higher risk of developing other conditions, such as ocular hypertension (OH), cataracts, cortisone-induced glaucoma, dry eye, diplopia and blindness [8,9], which in some cases can persist even after ceasing treatment with the drug. Among the chemotherapeutic drugs, Vincristine is associated with alterations of ocular motricity, corneal hypoesthesia, and optic atrophy [2]. Moreover, Methotrexate can induce alterations of motricity, such as internuclear ophthalmoplegia or optic neuropathy, with important repercussions for visual acuity [10]. Cytarabine, in turn, is associated with corneal toxicity [1].

This study characterized the ocular manifestations of pediatric patients during treatment for ALL and assessed whether such manifestations are associated with recurrence risk predictors defined in literature or with other factors, such as chemotherapeutic treatment protocols, gender, and infiltration of CSF. Early identification and treatment of such ocular alterations can prevent permanent visual loss, as well as precocious diagnosing of possible infiltration or incipient recurrence of the disease.

## **METHOD**

A prospective cohort study was conducted with children and adolescents diagnosed with ALL, receiving treatment between January 1<sup>st</sup>, 2013 through to December 30<sup>th</sup>, 2017 at the only public service specialized in pediatric oncology in the state of Sergipe, Northeast region of Brazil. The project was approved by the Ethics Commission at the Federal University of Sergipe (CEP-UFS) which deals with research involving human beings, report number 214.759.

The patients were classified at diagnosis into two groups according to their risk of recurrence: patients with a high risk of relapse (HR) were those who met one of the following criteria: age equal to or greater than nine years, white cell count greater than or equal to 50.000 cells/mm<sup>3</sup> at the time of diagnosis, and immunophenotyping compatible with T-cell ALL. The low risk group (LR) encompassed those patients who did not meet any of the above criteria [11].

The ALL treatment protocols, used in this study, were proposed in 1999 and updated in 2009 (ALL-99 and ALL-09) by the “Brazilian Group for Treatment of Acute Lymphoblastic Leukemia during Infancy” of the Brazilian Society of Pediatric Oncology. At the end of the first 28 days of treatment (D28), the ALL-99 protocol called for a total dosage of 1120 mg/m<sup>2</sup> of Prednisone for the LR group and 1680 mg/m<sup>2</sup> of Prednisone for the HR

group [12]. The ALL-09 protocol calls for 420 mg/m<sup>2</sup> of Prednisone and 126 mg/m<sup>2</sup> of Dexamethasone for the LR group and 1260 mg/m<sup>2</sup> of Prednisone for the HR group.

The ALL-99 protocol was routinely used with all patients admitted until December 31<sup>th</sup>, 2014, whereas the ALL-09 was used for patients admitted afterwards.

In this study, patients were included whose diagnosis of ALL was confirmed by immunophenotyping of bone marrow or peripheral blood and who met all of the following criteria: at least one eye examination in the six-month follow-up period beginning at diagnosis, age below 19 years, absence of previous chemotherapeutic treatment, absence of prior ocular pathology, no use of systemic GC in the six months prior to the diagnosis and existence of technical conditions allowing for the performance of eye examinations.

The eye examination was carried out by a single ophthalmologist (C.Q.M.) and consisted of exams before the start of treatment (D0), on the eighth day after admission (D8), at the end of the remission induction phase, which corresponds to the 28<sup>th</sup> day of treatment (D28), and at the end of the six first months of treatment (D6months). The exams on D0 and D8 were performed on the hospital bed due to the fragile health of these patients at diagnosis and consisted of measurement of IOP, evaluations of the anterior chamber with a portable slit-lamp biomicroscope and of the posterior chamber with direct ophthalmoscopy, without mydriasis. The D28 and D6months examinations were performed at an ophthalmologist's office, analyzing the anterior chamber by slit-lamp biomicroscopy and the posterior chamber by indirect ophthalmoscopy after mydriasis induction with 0.01% Tropicamide, assessment of visual acuity (VA), and measurement of IOP.

In all exams, IOP was measured using a Perkins applanation tonometer, under local anesthesia with 0.5% Proparacaine eyedrops and Fluoresceine tracing dye. Intraocular pressure equal or superior to 21mmHg was considered compatible with OH [13].

Visual acuity was assessed with an optotype auto projector, with and without mydriasis. The Snellen chart was used and VA < 20/40 was considered VL.


Patients diagnosed with any ocular pathology received appropriate ophthalmological treatment.

Data analysis was performed using the SPSS (*Statistical Package for Social Sciences*) software, version 24.0. Categorical variables were described as absolute and relative frequencies and compared using Fischer's exact test with Cramer V-test for strength and the Benjamini-Hochberg False Discovery Rate test for multiple comparison correction. The

continuous variables were described in terms of average and standard deviation. The binomial test was also used to compare proportions between non-parametric groups and the Pearson's correlation test was used to analyze the relationship between normally-distributed variables. P-values < 0.05 were considered statistically significant.

## RESULTS

Out of a total of 78 patients admitted to the service between 2013 and 2017, 67 met the inclusion criteria. However, 12 died before D8. Fifty-five patients had at least one eye examination between D8 and D6months. These patients were divided into two groups: those with normal examination results (n=37) and patients who presented some ocular manifestation (n=18). In turn, this last group was further divided into two groups depending on their visual acuity (Figure 1).



**Figure 1:** Flow chart showing the distribution of patients screened for possible ocular alterations in the cohort studied.



Sixty-seven patients with an average age of 9.10 years ( $SD = 5.436$ ) were included, presented in Table 1 by protocol, gender, age range, risk, immunophenotyping, deaths, cerebrospinal fluid (CSF) infiltration and ocular manifestations.

**Table 1. Profile of the patients included in the cohort**

| Variables             | N (%)     |
|-----------------------|-----------|
| Protocol              | 67        |
| ALL-99                | 35 (52.2) |
| ALL-09                | 32 (47.8) |
| Gender                | 67        |
| Feminine              | 30 (44.8) |
| Masculine             | 37 (55.2) |
| Age Range             | 67        |
| < 9 Years             | 36 (53.7) |
| ≥ 9 Years             | 31 (46.3) |
| Risk                  | 65        |
| Low                   | 25 (38.5) |
| High                  | 40 (61.5) |
| Immunophenotyping     | 67        |
| B-Cell                | 62 (92.5) |
| T-Cell                | 5 (7.5)   |
| Deaths                | 67        |
| Yes                   | 29 (43.3) |
| No                    | 38 (56.7) |
| CSF Infiltration      | 22        |
| Infiltrated           | 15 (68.2) |
| Negative              | 7 (31.8)  |
| Ocular Manifestations | 55        |
| Present               | 18 (32.7) |
| Absent                | 37 (67.3) |

The patients that presented ophthalmic alterations ( $N=18$ ) were mainly from the ALL-99 protocol (77.8%), male (55.6%), older than 9 years-old (55.6%), high-risk (83.3%), B-cell Immunophenotype (83.3%), had not died during the 6-month follow-up (55.6%), and 50% of them presented infiltration of CSF in at least one examination. Patients with ophthalmic alterations, when compared to the group with normal examination results, predominantly used

the ALL-99 protocol ( $p=0.022$ ) and belonged to the HR group ( $p=0.041$ ). These results were confirmed by the Benjamini-Hochberg test with a false discovery rate of 20% and both associations are strong, according to the Cramer V-test (0.32 and 0.34, respectively). These results can be seen in Table 2.

**Table 2.** Comparison of patient groups with normal and altered ophthalmic examination for different variables.

| Variables         | Group with Ophthalmic Alterations | Group with Normal Exam Results | P Value* | Corrected P-value** |
|-------------------|-----------------------------------|--------------------------------|----------|---------------------|
|                   | N (%)                             | N (%)                          |          |                     |
| Protocol          | 18                                | 37                             | 0.022    | 0.029               |
| ALL-99            | 14 (77.8)                         | 16 (43.2)                      |          |                     |
| ALL-09            | 4 (22.2)                          | 21 (56.8)                      |          |                     |
| Gender            | 18                                | 37                             | 1.000    | 0.200               |
| Feminine          | 8 (44.4)                          | 15 (40.5)                      |          |                     |
| Masculine         | 10 (55.6)                         | 22 (59.5)                      |          |                     |
| Age Range         | 18                                | 37                             | 0.566    | 0.143               |
| < 9 years         | 8 (44.4)                          | 21 (56.8)                      |          |                     |
| ≥ 9 years         | 10 (55.6)                         | 16 (43.2)                      |          |                     |
| Risk              | 17                                | 36                             | 0.041    | 0.057               |
| Low               | 3 (17.6)                          | 18 (50.0)                      |          |                     |
| High              | 14 (82.4)                         | 18 (50.0)                      |          |                     |
| Immunophenotyping | 18                                | 37                             | 0.671    | 0.171               |
| B-Cell            | 15 (83.3)                         | 33 (89.2)                      |          |                     |
| T-Cell            | 3 (16.7)                          | 4 (10.8)                       |          |                     |
| Deaths            | 18                                | 37                             | 0.213    | 0.086               |
| Yes               | 8 (44.4)                          | 9 (24.3)                       |          |                     |
| No                | 10 (55.6)                         | 28 (75.7)                      |          |                     |
| CSF Infiltration  | 4                                 | 16                             | 0.587    | 0.114               |
| Altered           | 2 (50.0)                          | 11 (68.8)                      |          |                     |
| Negative          | 2 (50.0)                          | 5 (31.3)                       |          |                     |

\* Fisher exact-test

\*\* Benjamini-Hochberg False Discovery Rate

Intra and/or extra-ocular alterations occurred in 32.7% of the patients examined, with a predominance of OH (61.1%), followed by retinal hemorrhage and VL (22.2%). Some

patients accumulated more than one alteration and the distribution of ophthalmic changes per patient is shown in Table 3.

**Table 3.** Distribution of eye manifestations per patient.

| <i>Patients</i>                                 | <i>1</i> | <i>2</i> | <i>3</i> | <i>4</i> | <i>5</i> | <i>6</i> | <i>7</i> | <i>8</i> | <i>9</i> | <i>10</i> | <i>11</i> | <i>12</i> | <i>13</i> | <i>14</i> | <i>15</i> | <i>16</i> | <i>17</i> | <i>18</i> |
|---|----------|----------|----------|----------|----------|----------|----------|----------|----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| <i>Alteration of retinal pigment epithelium</i> |          |          |          |          |          |          |          |          |          |           |           |           |           |           |           | x         |           |           |
| <i>Amaurosis</i>                                |          |          |          |          |          |          |          |          |          |           |           |           |           |           | x         |           |           |           |
| <i>Chorioretinal Atrophy</i>                    |          |          |          |          |          |          |          |          |          |           |           |           |           |           | x         |           |           |           |
| <i>Visual Loss</i>                              |          | x        |          |          |          |          |          |          |          |           |           |           |           |           | x         | x         | x         |           |
| <i>Orbital Cellulitis</i>                       |          |          | x        |          |          |          |          |          |          |           |           |           |           |           |           |           |           |           |
| <i>Papilla Edema</i>                            |          |          |          |          |          |          |          |          |          |           |           |           |           |           | x         |           |           |           |
| <i>Retinal Hemorrhage</i>                       |          |          |          | x        | x        |          |          |          |          |           |           |           |           |           | x         | x         |           |           |
| <i>Ocular Hypertension</i>                      |          |          |          |          |          |          | x        | x        | x        | x         | x         | x         | x         | x         |           | x         | x         | x         |
| <i>Leukemic Infiltration</i>                    |          |          |          |          |          |          |          |          |          |           |           |           |           |           |           |           |           | x         |
| <i>Optic Nerve Pallor</i>                       | x        | x        |          |          |          |          |          |          |          |           |           |           |           |           | x         |           |           |           |
| <i>3rd Nerve Palsy</i>                          |          |          |          |          |          |          |          |          |          |           |           |           |           |           | x         |           |           |           |
| <i>Retinochoroiditis</i>                        |          |          |          |          |          | x        |          |          |          |           |           |           |           |           |           |           |           |           |
| <i>Uveitis from Herpes</i>                      |          |          |          |          |          |          |          |          |          |           |           |           |           |           |           |           | x         |           |
| <i>Zoster</i>                                   |          |          |          |          |          |          |          |          |          |           |           |           |           |           |           |           |           |           |

When calculating the relative risk concerning the manifestation of ophthalmic changes using the cited variables, it was observed that the patients from the 1999 protocol have a risk 2.91 times higher for ocular manifestations than those from the 2009 protocol (CI 95% 1.099-


7.742) and low risk of relapse acting as a protective factor for presenting ocular manifestations (RR=0.327, CI 95% 0.107-0.999). These results are shown in Table 4.

**Table 4.** Relative risk of ocular manifestations.

|                  | RR    | CI 95%*     |
|------------------|-------|-------------|
| Deaths           | 1.788 | 0.859-3.722 |
| Immunophenotype  | 0.729 | 0.281-1.891 |
| CSF Infiltration | 0.538 | 0.095-3.041 |
| Protocol         | 2.917 | 1.099-7.742 |
| Gender           | 1.113 | 0.521-2.379 |
| Age Range        | 0.717 | 0.334-1.540 |
| Risk of relapse  | 0.327 | 0.107-0.999 |

\*95% confidence interval (CI)

The central tendency measurements with regard to IOP are shown using boxplots in Figure 2. It was observed that higher IOPs were found at D8, including some outliers with values of 24, 26 and 27 mmHg. The highest median was also observed at D8, approximately 17 mmHg. The lowest median, in turn, occurs at D6months, with a value of 14 mmHg. The greatest variation was observed at D28, whereas the smallest variations were observed to be similar at D0 and D6months.



**Figure 2:** Boxplots showing the distribution of Intraocular Pressure in patients during the six-month follow-up period.

## DISCUSSION

Childhood lymphoproliferative neoplasms most commonly attack B-cells and can affect the ocular appendix, including orbits, conjunctiva, eyelids and lacrimal glands [3]. Eye manifestations associated with the treatment of ALL are common, especially after the considerable increase in survival rate that followed the use of standardized multicentric protocols and better risk stratification, including the immunophenotyping of blasts. Furthermore, ocular impairment may be the first manifestation or sign of recurrence of the disease [13].

During the treatment, 32,7% of patients were confirmed as having ocular impairment, contrasting with previous studies that obtained 17%, 15%, 20% [3,14,16], but approaching others which found 29%, 39%, 38% [15, 17,18]. This data divergence is recurrent in the literature and reflects the application of different methods. Furthermore, in the present study, the highest proportion of ocular changes is due to the inclusion of the OH cases, which were actively traced in this cohort, even in the absence of symptoms, as was previously reported [9].

Risk of relapse acted as a protective factor, with a relative risk of 0.33, meaning that it is 67% less likely that one with low risk of relapse develop some ocular manifestation than one from the high-risk group, compatible with results in the literature [14]. This is also valid for those treated with the ALL-99 protocol, who had 2.91 times greater risk of presenting ocular manifestations than the ALL-09 patients, which could be explained by the lower total doses of GC in this group.

Direct ocular involvement occurred in one patient (2%) as leukemic infiltration of the optic nerve, an alteration observed in 7.3% of the cases in a previous study [15]. Among secondary ophthalmic alterations, the retina was the most affected tissue, with hemorrhage occurring in 7.3% of the cases, with similar proportions consistently described previously: 8% and 8.8% [3,16].

Other secondary alterations were the development of orbital cellulitis (2%) and uveitis due to Herpes zoster (2%). Patients with ALL can contract opportunistic germ infections and reactivate viral infections during the neutropenia phase, resulting in an elevated risk of death [19].

It was observed that during the remission induction phase, when high doses of GC are used, there was significant elevation of IOP in ten patients (20%), whose values were compatible with OH (IOP > 21mmHg). The result was greater than that presented in another study which reported 16.6% [9]. As in previous studies [3,9,13,14,15,19], no patient had developed isolated OH symptoms, making a diagnosis difficult when there is no systematic tracking of all patients.

Four patients suffered definitive visual impairment (7.3%) and none of those patients had developed isolated OH. These patients had other alterations secondary to the disease, such as retinal hemorrhage, optic nerve pallor, leukemic infiltration and uveitis, and suffered from reduced visual acuity.

There are remarkable differences observed between low and middle-income countries compared to that of developed countries in relation to indexes of mortality during induction, mortality at six months after diagnosis (the beginning of the maintenance phase), CSF infiltration and the proportion of patients with high risk of relapse [20]. These differences are more directly related to access to adequate infrastructure for treatment of complications than to the biological aspects of the disease. The data related to avoidable ocular complications, when promptly identified, can probably be generalized to apply to countries of all levels of income.

Thus, there exists the possibility of ocular involvement and silent OH, with the consequent risk of irreversible visual loss, in patients without any ocular alterations before the diagnosis. Thus, there is a need for systematic and universal ophthalmological evaluation of children and adolescents with neoplasia in the lymphoid lineage from the very beginning of treatment.

## CONCLUSION

Patients with ALL have elevated incidences of ocular manifestations due to the treatment and the disease itself. Among those, OH is the most prevalent. Patients treated with a protocol using higher doses of GC (ALL-99) and with high risk of recurrence had a higher risk of presenting ocular alterations, with strong association between those variables. Therefore, a protocol is proposed that contemplates systematic ophthalmological examination with the measurement of IOP immediately after diagnosis of ALL, before the introduction of GC (D0) and subsequently at D8, D28 and D6months.

## REFERENCES

- [1] Talcott KE, Garg RJ, Garg SJ. Ophthalmic manifestations of leukemia. *Current Opinion in Ophthalmology* 2016; **27**: 545–551.
- [2] Sharma T, Grewal J, Gupta S, Murray PI. Ophthalmic manifestations of acute leukaemias: the ophthalmologists role. *Eye* 2004; **18**: 663–672.
- [3] Reddy SC, Menon BS. A prospective study of ocular manifestations in childhood acute leukaemia. *Acta Ophthalmologica Scandinavica* 1998; **76**: 700–703.
- [4] Yalcinbayir O, Baytan B, Gelisken O, Can B, Evim MS, Yildiz M *et al.* Spectral domain optical coherence tomography findings of patients under treatment for pediatric acute lymphoblastic leukemia. *Journal of American Association for Pediatric Ophthalmology and Strabismus* 2017; **21**. doi:10.1016/j.jaapos.2016.12.002
- [5] Çaçı I, Ünlü K, Ari S, Dirier A. Unilateral Optic Disc Edema in a Patient With Acute Lymphocytic Leukemia: A Case and Review of the Literature. *Annals Of Ophthalmology* 2005; **37**: 303–306.
- [6] Chaudhuri T, Roy S, Roy P. Ischaemic optic neuropathy induced sudden blindness as an initial presentation of acute lymphoblastic leukemia. *Indian Journal of Medical and Paediatric Oncology* 2013; **34**: 335.
- [7] Gordon K. Ocular manifestations of leukemia leukemic infiltration versus infectious process. *Ophthalmology* 2001; **108**: 2293–2300.
- [8] Whelan KF, Stratton K, Kawashima T, Waterbor JW, Castleberry RP, Stovall M *et al.* Ocular late effects in childhood and adolescent cancer survivors: A report from the childhood cancer survivor study. *Pediatric Blood & Cancer* 2010; **54**: 103–109.
- [9] Mendonça CDQ, Souza CPD, Martins-Filho PRS, Viana SS, Leal BC, Cipolotti R. Steroid-induced ocular hypertensive response in children and adolescents with acute lymphoblastic leukemia and non-hodgkin lymphoma. *Pediatric Blood & Cancer* 2014; **61**: 2083–2085.
- [10] Fraunfelder F, Meyer SM. Ocular Toxicity of Antineoplastic Agents. *Ophthalmology* 1983; **90**: 1–3.

- [11] Cerdà-Ibáñez M, Bayo-Calduch P, Manfreda-Domínguez L, Duch-Samper A. Acute Vision Loss As The Only Sign Of Leukemia Relapse. *Retinal Cases & Brief Reports* 2018; **12**: 10–11.
- [12] Brandalise, S. R. et al. Benefits of the intermittent use of 6-mercaptopurine and methotrexate in maintenance treatment for low-risk acute lymphoblastic leukemia in children: randomized trial from the Brazilian Childhood Cooperative Group--protocol ALL-99. *Journal of Clinical Oncology*, v. 28, n. 11, p. 1911–1918, 10 abr. 2010.
- [13] Curto ML, Zingone A, Acquaviva A, Bagnulo S, Calculli L, Cristiani L *et al.* Leukemic infiltration of the eye: Results of therapy in a retrospective multicentric study. *Medical and Pediatric Oncology* 1989; **17**: 134–139.
- [14] Russo V, Scott I, Querques G, Stella A, Barone A, Noci ND. Orbital and Ocular Manifestations of Acute Childhood Leukemia: Clinical and Statistical Analysis of 180 Patients. *European Journal of Ophthalmology* 2008; **18**: 619–623.
- [15] Khadka D, Sharma A, Shrestha JK, Shrestha GS, Shrestha PN, Pant SR *et al.* Ocular manifestations of childhood acute leukemia in a tertiary level eye centre of Kathmandu, Nepal. *Nepalese Journal of Ophthalmology* 2014; **6**. doi:10.3126/nepjoph.v6i2.11678.
- [16] Bitirgen G, Belviranlı S, Caliskan U, Tokgoz H, Ozkagnici A, Zengin N. Ophthalmic Manifestations in Recently Diagnosed Childhood Leukemia. *European Journal of Ophthalmology* 2015; **26**: 88–91.
- [17] Reddy S, Jackson N, Menon B. Ocular Involvement in Leukemia – A Study of 288 Cases. *Ophthalmologica* 2003; **217**: 441–445.
- [18] Schachat AP. Ophthalmic Manifestations of Leukemia. *Archives of Ophthalmology* 1989; **107**: 697.
- [19] Cogan DG. Immunosuppression and Eye Disease. *American Journal of Ophthalmology* 1977; **83**: 777–788.
- [20] Howard SC, Metzger ML, Wilimas JA, Quintana Y, Pui CH, Robison LL, Ribeiro RC. Childhood Cancer Epidemiology in Low-Income Countries. *Cancer* 2008 Feb 1; **112**(3):461-72. Review.



## **5.2 Estudo de Revisão Sistemática (Artigo 2)**

### **Steroid-Induced Ocular Hypertensive Response in Children and Adolescents with Acute Lymphoblastic Leukemia and Non-Hodgkin Lymphoma: a Systematic Review**

(Artigo submetido no “Journal of Pediatric Hematology and Oncology JPHO-18-150. Foram submetidas as correções solicitadas pelos revisores; manuscrito com o Editor, aguardando parecer final).

#### **Summary**

The aim of this study was to conduct a systematic review to evaluate the risk of increased intraocular pressure (IOP) related to corticosteroids in children and adolescents with acute lymphoblastic leukemia (ALL) and non-Hodgkin lymphoma (NHL). Eligible studies had to include children and adolescents up to 19 years of age with ALL or NHL, regardless of gender, and treated with systemic corticosteroids on a regular schedule. Four studies met the eligibility criteria: two were individual case reports, one a series of five cases and one observational prospective study of twelve patients. The results described eventual control of IOP, and outcomes ranging from no eye damage to total blindness. The possibility of silent

ocular hypertension, with consequent risk of irreversible blindness, indicates the need for evaluating the introduction of an IOP verification protocol in young patients immediately after diagnosis of ALL or NHL.

However, due to limited evidence, further studies on evaluation of IOP and visual function before starting treatment, as well as systematic measurements during and after treatment, are warranted.

**Key words:** ocular hypertension, steroids, acute lymphoblastic leukemia.

## **Introduction**

Acute lymphoblastic leukemia (ALL) is the most common cancer during childhood, corresponding to 32% of neoplasms and 85% of all cases of childhood leukemia.<sup>1</sup> Non-Hodgkin lymphoma (NHL) accounts for approximately 4% of all malignancies among children.<sup>2</sup>

The first description of corticosteroid-induced ocular hypertension was published in 1950.<sup>3</sup> The disorder occurs independent of the method of drug administration and constitutes an important and preventable cause of visual loss. Previous studies had already shown that the risk of ocular hypertension is higher in children<sup>4</sup> especially after some weeks of corticosteroid administration, and that this is generally reversible with the suspension of the drug's use. However, if the ocular hypertension persists for a prolonged period, permanent damage of the optical nerve may occur, with consequent vision impairment.<sup>5</sup>

The aim of this study was to conduct a systematic review to evaluate the risk of increased intraocular pressure (IOP) and its possible implications for visual function related to corticosteroid therapy in children and adolescents with ALL and NHL.

## **Methods**

A protocol of this systematic review was designed a priori and was registered in the PROSPERO database (registration number CRD42018087146).

## **Literature Search**

We searched PubMed, Scopus, Web of Science, Science Direct, and Cochrane Library from inception to January 31, 2018. A grey-literature search was conducted through Google Scholar and OpenThesis. The first 100 results of Google Scholar were recorded. The search also included a hand search of cross-references from original articles and reviews. The structured search strategy used the following terms: (precursor cell lymphoblastic leukemia-

lymphoma OR acute lymphoblastic leukemia OR non-Hodgkin lymphoma) AND (glaucoma OR ocular hypertension). Two independent reviewers screened the search results and identified studies that were potentially relevant based on article titles and abstracts.

Relevant studies were read in full and selected according to the eligibility criteria.

### **Eligibility Criteria**

In this systematic review, we included studies that reported the occurrence of corticosteroid-induced ocular hypertension in children and adolescents with ALL or NHL. Eligible studies had to include children and adolescents from 5 to 19 years of age, regardless of gender, and treated with systemic corticosteroids on a regular schedule for ALL or NHL. If the assessed study included several types of malignancies treated with systemic corticosteroid, separate data for ALL and NHL had to have been provided. Studies reporting ocular hypertension by tumoral infiltration, and glaucomatous field damage and outcome of corticosteroid-induced ocular hypertension at initial testing were excluded. We did not exclude any study on criteria of sample size, duration of follow-up, publication date, or language of publication.

### **Data Extraction**

Two reviewers independently abstracted data from studies that met the eligibility criteria using a predefined coding protocol. Disagreements were resolved by consensus or, rarely, adjudication by a third reviewer. We extracted information on (1) design and demographic details, including type of study, number of patients, geographic location, type of malignancy, gender, age, treatment (chemotherapy, radiotherapy, or systemic corticosteroid use), previous ocular pathology, family history of glaucoma, diagnosis of corticosteroid-induced ocular hypertension, and standards for measurement of ocular pressure; (2) details of corticosteroid use and IOP, including data on induction and post-induction therapy (type of corticosteroid, course, and dosage), cycles, baseline IOP and values of ocular hypertension; (3) strategies to control IOP, including use of antiglaucoma medication, corticosteroid switching, corticosteroid cessation, and complications.

### **Exposure and Outcome**

The exposure was defined as treatment of ALL or NHL with systemic corticosteroids, regardless of type of corticosteroid or dosage. The outcome was the ocular hypertension occurring after the initiation of corticosteroid and defined as an IOP higher than 21 mmHg in one or both eyes.

## Results

Altogether, 314 reports were retrieved and, after reviewing the titles and abstracts, 9 full-text articles were assessed for eligibility. Of these, 5 were excluded.<sup>6-10</sup> The remaining 4 articles<sup>11-14</sup> met the eligibility criteria and were included in the systematic review (Figure 1). The 4 reviewed articles<sup>11-14</sup> were a case report in the form of a summary<sup>11</sup>, published for the American Society of Pediatric Hematology/Oncology 25th Annual Meeting, 2012, New Orleans, Louisiana; and 3 complete studies, designed as a case report<sup>12</sup>, a series of 5 cases<sup>13</sup> and an observational prospective study with 12 patients.<sup>14</sup> All included studies were published in the English language and the outcome under evaluation was the presence of ocular hypertension. No studies were multicenter or randomized.

The first study was presented as a poster at the American Society of Pediatric Hematology/Oncology 25th Annual Meeting, New Orleans, Louisiana, in 2012, by Pilbeam et al.<sup>11</sup> The authors reported the case of a 3-year-old boy diagnosed with ALL, who had no previous history of visual problems or family history of glaucoma. On the 10th day of induction with dexamethasone 3 mg/m<sup>2</sup>, the patient presented with photophobia, irritability, decreased visual acuity and dilated pupils. The initial diagnostic hypothesis was optic neuropathy, but the brain magnetic resonance and angioresonance were normal. Ocular exam showed visual acuity at 20/200, dilated non-reactive pupils and elevated IOP (OD 42 mmHg, OS 40 mmHg). The patient, although treated for glaucoma, developed permanent blindness in the right eye and a significant loss of vision in the left eye.

In the second study, Tham et al.<sup>12</sup> reported the case of a 9-year-old-girl with ALL who received high doses of systemic corticosteroid (60 mg daily of oral prednisolone) and presented on the 10th day of treatment an IOP of 52 mmHg in the right eye and 47 mmHg in the left eye. Topical medications to treat glaucoma were used (latanoprost 0.001% once daily and brimonidine 0.2% twice daily) and the pressure dropped to 38 mmHg and 36 mmHg. IOP returned to baseline after 2 days of suspension of the corticosteroid and after 6 weeks of follow-up, this level was maintained even without glaucoma

medications. Four months later, the patient was submitted to a new cycle of treatment with 10 mg of oral dexamethasone daily. The patient showed similar increase in IOP for both eyes

during the course of oral dexamethasone. Oral acetazolamide was prescribed and the patient remained asymptomatic throughout treatment, except for one episode of decreased visual acuity in the right eye when IOP reached 52 mmHg.

In the third study, Yamashita et al.<sup>13</sup> reported a case series including children up to 6 years of age diagnosed with ALL (2 boys and 3 girls, mean  $4.2 \pm 1.6$  years). The 15 treatment cycles were each composed of induction therapy (prednisolone 60 mg/m<sup>2</sup>/day for 4 weeks) followed by late intensification therapy (dexamethasone 6mg/m<sup>2</sup>/day for 2 weeks) and a maintenance regime (dexamethasone, 6mg/m<sup>2</sup>/day for 2 weeks, then 1 week of tapering and a 5-week break). The patients with ocular symptoms were referred to an ophthalmologist throughout the treatment. Children were examined for visual acuity and IOP was measured by non-contact tonometry (CT-80, Topcon, Japan) on each 7th day of induction therapy and 7th day of maintenance regime. All patients had IOP above 21 mmHg, with a mean of  $39.6 \pm 7.2$  mmHg (range 28 to 47). Antiglaucoma medication was given to all presenting IOP above 25 mmHg. Of the 5 patients included in this study, one received only antiglaucoma medication; 3 were treated with antiglaucoma medication and the steroid switched from dexamethasone to prednisolone; and one showed a well-controlled IOP with antiglaucoma medication and cessation of corticosteroid therapy, even though he presented severe glaucomatous optic atrophy. In all cases, regardless of the method of IOP control, the pressure was normalized.

In the most recent study included in this review, Mendonca et al.<sup>14</sup> performed a prospective observational study with measurement of IOP before treatment (D0), and on the 8th (D8), 14th (D14), and 28th days (D28) of treatment. The authors examined 12 patients (mean age of  $8.8 \pm 5.4$  years), 11 diagnosed with ALL and one with NHL. Eight patients were given prednisone and 4 received dexamethasone. Two cases (16.7%) of ocular hypertension were observed, the first induced by dexamethasone with IOP at 42 mmHg in both eyes on D14, and the second case induced by prednisone with IOP at 26 mmHg in both eyes on D8. Satisfactory control of the IOP was obtained using eye drops of Brinzolamide 1.0% and timolol maleate 0.5 % twice a day. No patient presented ocular hypertension after D28, when the corticosteroids were suspended. Differences of IOP between D0 vs. D8 and D0 vs. D14 ( $p=0.013$ ) were observed. The results of this review are summarized in Tables 1 and 2.

## Discussion

This systematic review included 9 children who were diagnosed with ALL and presented steroid-induced ocular hypertension. No cases in children with NHL were reported.

Unfortunately, although we used an explicit method to identify and select relevant studies, a limited number were found for inclusion.

Only one study among those 4 that met the eligibility criteria had been prospective<sup>14</sup> and had estimated the incidence of the steroid-induced ocular hypertension in a sample of 12 children. The increased risk of steroid-induced ocular hypertension remains controversial. Recently, Whelan et al<sup>6</sup> in a multi-institutional collaborative study of individuals who survived at least 5 years after diagnosis of cancer during childhood or adolescence, did not find an increased risk of glaucoma among survivors treated with glucocorticoid therapy. However, the authors suggest that the true incidence of ocular hypertension may be underestimated, since patients can remain asymptomatic, that regular eye exams and additional follow-up may uncover increased risk of glaucoma in childhood cancer survivors treated with corticosteroids and that it is also important that IOP be measured at the beginning of corticosteroid therapy, making it possible to evaluate the real increase in pressure levels and the ideal reading to indicate their control. Of the 4 studies included in this review, only 2 studies<sup>12,14</sup> reported the measurement of ocular pressure at the beginning of corticosteroid treatment.

The use of varied methods to measure IOP can also explain differences in the incidence of ocular hypertension in children receiving corticosteroid therapy. In the study performed by Yamashita et al.<sup>13</sup>, IOP was measured by air puff, non-contact tonometry (Topcon CT-80) that does not have the reliability of contact tonometry, resulting in possible overestimation of values in children. Pediatric patients with ALL or NHL receiving corticosteroid can remain asymptomatic in spite of severe ocular hypertension.

Thus, controlled studies with evaluation of IOP by applanation tonometry become necessary in order to insure more precise diagnoses and treatment decisions, as shown by Mendonca et al.<sup>14</sup>

All patients diagnosed with steroid-induced ocular hypertension included in this study were treated with ocular hypotensive medications with adequate control of IOP in most cases. However, some serious complications were observed in the studies by Pilbeam et al.<sup>11</sup>, Tham et al.<sup>12</sup>, and Yamashita et al.<sup>13</sup>, including severe glaucomatous optic atrophy and permanent blindness. According to Tham et al.<sup>12</sup>, the most effective way to normalize the IOP appeared to be cessation of the corticosteroid therapy, but this conduct is impossible in most cases due to the nature of the underlying medical condition. In the study performed by Yamashita et al.<sup>11</sup>, different methods were used to control the IOP, but all of them included the use of antiglaucoma medications. The use of ocular hypotensive medications thus appears to be an


adequate choice for the control of steroid-induced ocular hypertension in children, whether the type of steroid is switched or not. However, further studies are needed to verify the best combination therapy in the treatment of steroid-induced ocular hypertension in pediatric patients with ALL and NHL.

In this review, only the study performed by Mendonca et al.<sup>14</sup> showed measurements of IOP and visual function at the beginning of corticoid induction, as well as systematic measurements during and after the treatment. The authors observed a significant increase of IOP compared to baseline during the remission induction phase of treatment, when elevated doses of glucocorticosteroids were used. In 2 out of 12 patients (16.7%), the IOP levels could have been high enough to cause glaucomatous optic neuropathy and eventually blindness, since the risk of glaucomatous nerve damage is directly related to the level and duration of IOP.<sup>11,15</sup>

## **Conclusion**

The possibility of silent ocular hypertension, with consequent risk of irreversible blindness, indicates the need for evaluating the introduction of an IOP verification protocol in young patients immediately after diagnosis of ALL or NHL. However, due to limited evidence, further studies on evaluation of IOP and visual function before starting treatment, as well as systematic measurements during and after treatment, are warranted.

Figure 1 **Flow diagram.**



**Table 1. Design features and demographic details of studies included in the systematic review.**

| Study                          | Type of study             | Number of patients | Country | Malignancy  | Sex    | Age         | Treatment |    |         | SIOH         | Measurement                          |
|--------------------------------|---------------------------|--------------------|---------|-------------|--------|-------------|-----------|----|---------|--------------|--------------------------------------|
|                                |                           |                    |         |             |        |             | CT        | RT | Oral GC |              |                                      |
| Mendonca et al. <sup>14</sup>  | Prospective observational | 12                 | Brazil  | ALL and NHL | 5F:7 M | 8.8y (mean) | Yes       | No | Yes     | 2 patients   | Perkins applanation tonometry        |
| Yamashita et al. <sup>13</sup> | Case series               | 5                  | Japan   | ALL         | 3F:2 M | 4.2y (mean) | Yes       | No | Yes     | All patients | Noncontact tonometry (CT-80, TOPCON) |
| Tham et al. <sup>12</sup>      | Case report               | 1                  | China   | ALL         | F      | 9y          | ND        | ND | Yes     | Yes          | Goldmann applanation tonometry       |
| Pilbeam et al. <sup>11</sup>   | Case report               | 1                  | USA     | ALL         | M      | 3y          | Yes       | No | Yes     | Yes          | ND                                   |

CT: chemotherapy; RT: radiotherapy; ND: not described; SIOH: steroid-induced ocular hypertension.

**Table 2. IOP control and complications in patients with steroid-induced ocular hypertension.**



| Study                          | Patient | Method                  |                   |                   | IOP               | Complications  |
|--------------------------------|---------|-------------------------|-------------------|-------------------|-------------------|--|
|                                |         | Antiglaucoma medication | Steroid switching | Steroid cessation |                   |  |
| Mendonca et al. <sup>14</sup>  | 1       | Yes                     | No                | No                | Well controlled   | No   |
|                                | 2       | Yes                     | No                | No                | Well controlled   | No   |
| Yamashita et al. <sup>13</sup> | 3       | Yes                     | Yes               | No                | Well controlled   | No   |
|                                | 4       | Yes                     | No                | No                | Well controlled   | No   |
|                                | 5       | Yes                     | Yes               | No                | Well controlled   | No   |
|                                | 6       | Yes                     | No                | Yes               | Well controlled   | Severe glaucomatous optic atrophy and decrease of retinal nerve fibers layer thickness |
|                                | 7       | Yes                     | Yes               | No                | Well controlled   | No   |
| Tham et al. <sup>12</sup>      | 8       | Yes                     | No                | No                | Poorly controlled | One episode of decreased visual acuity   |
| Pilbeam et al. <sup>11</sup>   | 9       | Yes                     | No                | No                | Poorly controlled | Permanent blindness in the right eye and significant vision loss in the left eye       |

## References

1. Yasmeen N, Ashraf S. Childhood acute lymphoblastic leukaemia; epidemiology and clinicopathological features. *J Pak Med Assoc.* 2009;59(3):150–153.
2. Alexander DD, Mink PJ, Adami HO, et al. The non-Hodgkin lymphomas: A review of the epidemiologic literature. *Int J Cancer.* 2007;120(S12):1–39.
3. McLean JM. Use of ACTH and cortisone. *Trans Am Ophthalmol Soc.* 1950; 48:293-296.
4. Brito P, Silva S, Cotta J, Falcão-Reis F. Severe ocular hypertension secondary to steroid treatment in a child with nephrotic syndrome. *Clin Ophthalmol.* 2012; 6:1675-9.
5. Danias J, Gerometta R, Ge Y, et al. Gene expression changes in steroid-induced IOP elevation in bovine trabecular meshwork. *Invest Ophthalmol Vis Sci.* 2011;52(12):8636–8645.
6. Whelan KF, Stratton K, Kawashima T, et al. Ocular late effects in childhood and adolescent cancer survivors: a report from the childhood cancer survivor study. *Pediatr Blood Cancer.* 2010;54(1):103–109.

7. Yamamoto-Ogasawara A, Asakawa M, Yoshino K, Nagamoto T, Inoue M, Hirakata A. Anterior complications in case of recurrent acute lymphoblastic leukaemia diagnosed by biopsy of aqueous humour. *Clin Exp Ophthalmol*. 2009;37(7):739–741.
8. Huang LT, Hsiao CC, Weng HH, Lui CC. Neurologic complications of pediatric systemic malignancies. *J Formos Med Assoc*. 1996;95(3):209–212.
9. Kurup SK, Levy-Clarke G, Calvo KR, Jaffe ES, Nussenblatt RB, Chan C-C. Primary diffuse large B-cell lymphoma of the spleen with coincident serous retinal detachments responsive to corticosteroids. *Clin Exp Ophthalmol*. 2007;35(5):468–472.
10. Cristol SM, Baumbblatt JG, Icasiano E, Grajewski AL, Olivier MMG, Parrish RK. Bilateral acute angle-closure associated with systemic lymphoma: a report of 2 cases. *J Glaucoma*. 2011;20(2):115–117.
11. Pilbeam K, Salvi S, Havani A. Corticosteroid-induced glaucoma as a complication of induction therapy in a child with acute lymphoblastic leukemia (ALL). *ASPHO Abstracts. Pediatr Blood Cancer*. 2012; 58:1014-1097.
12. Tham C, Ng J, Li R, Chik KW. IOP profile of a child on a systemic corticosteroid. *Am J Ophthalmol*. 2004; 137(1): 198-201.
13. Yamashita T, Kodama Y, Tanaka M, Yamakiri K, Kawano Y, Sakamoto T. Steroid-induced glaucoma in children with acute lymphoblastic leukemia: a possible complication. *J Glaucoma*. 2010;19(3):188–190.
14. Mendonca C, de Souza CP, Martins-Filho PRS, Viana SS, Leal BC, Cipolotti R. Steroid-induced ocular hypertensive response in children and adolescents with acute lymphoblastic leukemia and non-Hodgkin lymphoma. *Pediatr Blood Cancer*. 2014;61(11):2083–2085.
15. Bresson-Dumont H. IOP in children. *J Fr Ophtalmol*. 2009; Mar.32(3):176-81

### **5.3 Regras para Publicação**

#### **5.3.1 Estudo de Coorte (Artigo 1)**

#### **Description**

#### **Author Information Pack**

ISSN: 0145-2126

*Leukemia Research* an international journal which brings comprehensive and current information to all health care professionals involved in basic and applied clinical research in hematological malignancies. The editors encourage the submission of articles relevant to hematological malignancies. The Journal scope includes reporting studies of cellular and molecular biology, genetics, immunology, epidemiology, clinical evaluation, and therapy of these diseases.

### **Benefits To Authors**

We also provide many author benefits, such as free PDFs, a liberal copyright policy, special discounts on Elsevier publications and much more. Please [click here](#) for more information on our author services.

Please see our [Guide for Authors](#) for information on article submission. If you require any further information or help, please visit our [Support Center](#).

### **Audience**

Medical scientists, haematologists, immunologists, oncologists and chemotherapists.

### **Impact Factor**

2017: 2.319 © Clarivate Analytics Journal Citation Reports 2018

### **Abstracting And Indexing**

Chemica

Embase

Medline®

Elsevier Sociedad Iberoamericana de Informacion Cientifica (SIIC)

Pascal/Cnrs

Scopus

Science Citation Index Biosis

### **Editorial Board**

#### *Editor-in-Chief*

Clive S. Zent, James Wilmot Cancer Center, University of Rochester Medical Center, 601 Elmwood Avenue, Box 704, Rochester, New York, 14642, USA

## GUIDE FOR AUTHORS

### *Your Paper Your Way*

We now differentiate between the requirements for new and revised submissions. You may choose to submit your manuscript as a single Word or PDF file to be used in the refereeing process. Only when your paper is at the revision stage, will you be requested to put your paper in to a 'correct format' for acceptance and provide the items required for the publication of your article.

**To find out more, please visit the Preparation section below.**

### **Introduction**

*Leukemia Research* an international journal which brings comprehensive and current information to all health care professionals involved in basic and applied clinical research in hematological malignancies. The editors encourage the submission of articles relevant to hematological malignancies. The Journal scope includes reporting studies of cellular and molecular biology, genetics, immunology, epidemiology, clinical evaluation, and therapy of these diseases.

Word count limit does not include the abstract references, acknowledgement document, figures and tables.

**Original Articles** are full-length research papers which have not been published previously, except in a preliminary form. An Original paper usually does not exceed 4,000 words, it should be divided into sections (Introduction, Materials and Methods, Results and Discussion), 250 word abstract, longer papers may be considered if relevant, limit of 50 references and limit of 6 figures and/or tables.

**Review Articles** should be submitted after prior consultation with an Editor. This paper should not exceed 5,000 words but longer papers may be considered if relevant, limit of 150 references and no limit for figures and/or tables.

**Editorials** do not have abstracts and should not exceed 1,000 words, limit of 5 references and optional 1-2 figures and/or tables. Editorials should be submitted after prior consultation with an Editor.

**Correspondence**, which encompasses preliminary research observations and research letters, as well as conventional Letters to the Editor. No abstract required. 1,500 words (excluding references only) max 2 figures and tables, 20 References.

**Commentaries** are contributions invited by the Editor which should be concerned with matters of opinion and criticism on contributions published in the journal and other matters of interest to researchers in our field, often invited by the Editor to present new perspectives and advance understanding of controversial issues by provoking debate and comment. (1,500 words, 10 references, 1 figure, 1 table)

Authors may send queries concerning the submission process, manuscript status, or journal procedures to the Editorial Office(e-mail: LR@elsevier.com).

### ***Submission checklist***

You can use this list to carry out a final check of your submission before you send it to the journal for review. Please check the relevant section in this Guide for Authors for more details.

Author information pack: [www.elsevier.com/locate/leukres](http://www.elsevier.com/locate/leukres) 4

- Indicate clearly if color should be used for any figures in print *Graphical Abstracts* / *Highlights files* (where applicable) *Supplemental files* (where applicable). Further considerations
- Manuscript has been 'spell checked' and 'grammar checked'
- All references mentioned in the Reference List are cited in the text, and vice versa
- Permission has been obtained for use of copyrighted material from other sources.
- A competing interests statement is provided.
- Journal policies detailed in this guide have been reviewed
- Referee suggestions and contact details provided, based on journal requirements

For further information, visit our Support Center.

The Journal allows up to 6 months for authors to make the suggested revisions to their manuscript.

Note: If you do not submit your revisions by the 6 month date your paper will be withdrawn.

## **BEFORE YOU BEGIN**

### ***Ethics in publishing***

Please see our information pages on Ethics in publishing and Ethical guidelines for journal publication.

### ***Studies in humans and animals***

If the work involves the use of human subjects, the author should ensure that the work described has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans. The manuscript should be in line with the Recommendations for the Conduct, Reporting, Editing and Publication of Scholarly Work in Medical Journals and aim for the inclusion of representative human populations (sex, age and ethnicity) as per those recommendations. The terms sex and gender should be used correctly.

Authors should include a statement in the manuscript that informed consent was obtained for experimentation with human subjects. The privacy rights of human subjects must always be observed.

All animal experiments should comply with the ARRIVE guidelines and should be carried out in accordance with the U.K. Animals (Scientific Procedures) Act, 1986 and associated guidelines, EU Directive 2010/63/EU for animal experiments, or the National Institutes of Health guide for the care and use of Laboratory animals (NIH Publications No. 8023, revised 1978) and the authors should clearly indicate in the manuscript that such guidelines have been followed. The sex of animals must be indicated, and where appropriate, the influence (or association) of sex on the results of the study.

### ***Declaration of interest***

All authors must disclose any financial and personal relationships with other people or organizations that could inappropriately influence (bias) their work. Examples of potential competing interests include employment, consultancies, stock ownership, honoraria, paid expert testimony, patent applications/registrations, and grants or other funding. Authors must disclose any interests in two places: 1. A summary declaration of interest statement in the title page file (if double-blind) or the manuscript file (if single-blind). If there are no interests to declare then please state this: 'Declarations of interest: none'. This summary statement will be

ultimately published if the article is accepted. 2. Detailed disclosures as part of a separate Declaration of Interest form, which forms part of the journal's official records. It is important for potential interests to be declared in both places and that the information matches. More information.

### ***Submission declaration and verification***

Submission of an article implies that the work described has not been published previously (except in the form of an abstract, a published lecture or academic thesis, see 'Multiple, redundant or concurrent publication' for more information), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright- holder. To verify originality, your article may be checked by the originality detection service Crossref Similarity Check.

Please note that preprints can be shared anywhere at any time, in line with Elsevier's sharing policy. Sharing your preprints e.g. on a preprint server will not count as prior publication (see 'Multiple, redundant or concurrent publication' for more information).

### ***Use of inclusive language***

Inclusive language acknowledges diversity, conveys respect to all people, is sensitive to differences, and promotes equal opportunities. Articles should make no assumptions about the beliefs or commitments of any reader, should contain nothing which might imply that one individual is superior to another on the grounds of race, sex, culture or any other characteristic, and should use inclusive language throughout. Authors should ensure that writing is free from bias, for instance by using 'he or she', 'his/her' instead of 'he' or 'his', and by making use of job titles that are free of stereotyping (e.g. 'chairperson' instead of 'chairman' and 'flight attendant' instead of 'stewardess').

### ***Authorship***

All authors should have made substantial contributions to all of the following: (1) the conception and design of the study, or acquisition of data, or analysis and interpretation of data, (2) drafting the article or revising it critically for important intellectual content, (3) final approval of the version to be submitted.

### ***Role of Corresponding Author***

Please note: there is only one corresponding author of the manuscript. The corresponding author is the one that submits the manuscript and has the duty to ensure that all of the named authors have seen and approved the original manuscript and subsequent revisions. Each author should have participated sufficiently in the work to take public responsibility for the content. The corresponding author should also ensure that those who have contributed to the research are acknowledged appropriately either as a co-author or in the Acknowledgements. In addition, the corresponding author has the prime responsibility for ensuring the paper is correctly prepared according to the Guide for Authors. Submitted manuscripts not complying with the Guide for Authors may be returned to the authors.

### ***Changes to authorship***

Authors are expected to consider carefully the list and order of authors before submitting their manuscript and provide the definitive list of authors at the time of the original submission. Any addition, deletion or rearrangement of author names in the authorship list should be made only **before** the manuscript has been accepted and only if approved by the journal Editor. To request such a change, the Editor must receive the following from the corresponding author: (a) the reason for the change in author list and (b) written confirmation (e-mail, letter) from all authors that they agree with the addition, removal or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author being added or removed. Only in exceptional circumstances will the Editor consider the addition, deletion or rearrangement of authors after the manuscript has been accepted. While the Editor considers the request, publication of the manuscript will be suspended. If the manuscript has already been published in an online issue, any requests approved by the Editor will result in a corrigendum.

### **Clinical trial results**

In line with the position of the International Committee of Medical Journal Editors, the journal will not consider results posted in the same clinical trials registry in which primary registration resides to be prior publication if the results posted are presented in the form of a brief structured (less than 500 words) abstract or table. However, divulging results in other circumstances (e.g., investors' meetings) is discouraged and may jeopardise consideration of the manuscript. Authors should fully disclose all posting in registries of results of the same or closely related work.

#### *Reporting clinical trials*

Randomized controlled trials should be presented according to the CONSORT guidelines. At manuscript submission, authors must provide the CONSORT checklist accompanied by a flow diagram that illustrates the progress of patients through the trial, including recruitment, enrollment, randomization, withdrawal and completion, and a detailed description of the randomization procedure. The CONSORT checklist and template flow diagram are available online.

### **Registration of clinical trials**

Registration in a public trials registry is a condition for publication of clinical trials in this journal in accordance with International Committee of Medical Journal Editors recommendations. Trials must register at or before the onset of patient enrolment. The clinical trial registration number should be included at the end of the abstract of the article. A clinical trial is defined as any research study that prospectively assigns human participants or groups of humans to one or more health-related interventions to evaluate the effects of health outcomes. Health-related interventions include any intervention used to modify a biomedical or health-related outcome (for example drugs, surgical procedures, devices, behavioral treatments, dietary interventions, and process-of-care changes). Health outcomes include any biomedical or health-related measures obtained in patients or participants, including pharmacokinetic measures and adverse events. Purely observational studies (those in which the assignment of the medical intervention is not at the discretion of the investigator) will not require registration.

### **Article transfer service**

This journal is part of our Article Transfer Service. This means that if the Editor feels your article is more suitable in one of our other participating journals, then you may be asked to consider transferring the article to one of those. If you agree, your article will be transferred automatically on your behalf with no need to reformat. Please note that your article will be reviewed again by the new journal. More information.

## **Copyright**

Upon acceptance of an article, authors will be asked to complete a 'Journal Publishing Agreement' (see more information on this). An e-mail will be sent to the corresponding author confirming receipt of the manuscript together with a 'Journal Publishing Agreement' form or a link to the online version of this agreement.

Subscribers may reproduce tables of contents or prepare lists of articles including abstracts for internal circulation within their institutions. Permission of the Publisher is required for resale or distribution outside the institution and for all other derivative works, including compilations and translations. If excerpts from other copyrighted works are included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has preprinted forms for use by authors in these cases.

For gold open access articles: Upon acceptance of an article, authors will be asked to complete an 'Exclusive License Agreement' (more information). Permitted third party reuse of gold open access articles is determined by the author's choice of user license.

## **Author rights**

As an author you (or your employer or institution) have certain rights to reuse your work. More information.

## **Elsevier supports responsible sharing**

Find out how you can share your research published in Elsevier journals.

## **Role of the funding source**

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. If the funding source(s) had no such involvement then this should be stated.

### *Funding body agreements and policies*

Elsevier has established a number of agreements with funding bodies which allow authors to comply with their funder's open access policies. Some funding bodies will reimburse the author for the gold open access publication fee. Details of existing agreements are available online.

After acceptance, open access papers will be published under a noncommercial license. For authors requiring a commercial CC BY license, you can apply after your manuscript is accepted for publication.

## **Informed consent and patient details**



Studies on patients or volunteers require ethics committee approval and informed consent, which should be documented in the paper. Appropriate consents, permissions and releases must be obtained where an author wishes to include case details or other personal information or images of patients and any other individuals in an Elsevier publication. Written consents must be retained by the author but copies should not be provided to the journal. Only if specifically requested by the journal in exceptional circumstances (for example if a legal issue arises) the author must provide copies of the consents or evidence that such consents have been obtained. For more information, please review the Elsevier Policy on the Use of Images or Personal Information of Patients or other Individuals. Unless you have written permission from the patient (or, where applicable, the next of kin), the personal details of any patient included in any part of the article and in any supplementary materials (including all illustrations and videos) must be removed before submission.

## **Submission**

Our online submission system guides you stepwise through the process of entering your article details and uploading your files. The system converts your article files to a single PDF file used in the peer-review process. Editable files (e.g., Word, LaTeX) are required to typeset your article for final publication. All correspondence, including notification of the Editor's decision and requests for revision, is sent by e-mail.

### *Submit your article*

Please submit your article via <http://ees.elsevier.com/lr/>.

### *Referees*

Please submit the names and institutional e-mail addresses of several potential referees. For more details, visit our Support site. Note that the editor retains the sole right to decide whether or not the suggested reviewers are used.

## **Preparation**

### **New Submissions**

Submission to this journal proceeds totally online and you will be guided stepwise through the creation and uploading of your files. The system automatically converts your files to a single PDF file, which is used in the peer-review process.

As part of the Your Paper Your Way service, you may choose to submit your manuscript as a single file to be used in the refereeing process. This can be a PDF file or a Word document, in any format or lay-out that can be used by referees to evaluate your manuscript. It should contain high enough quality figures for refereeing. If you prefer to do so, you may still provide all or some of the source files at the initial submission. Please note that individual figure files larger than 10 MB must be uploaded separately.

### *References*

There are no strict requirements on reference formatting at submission. References can be in any style or format as long as the style is consistent. Where applicable, author(s) name(s), journal title/ book title, chapter title/article title, year of publication, volume number/book chapter and the article number or pagination must be present. Use of DOI is highly encouraged. The reference style used by the journal will be applied to the accepted article by Elsevier at the proof stage. Note that missing data will be highlighted at proof stage for the author to correct.

### *Formatting requirements*

There are no strict formatting requirements but all manuscripts must contain the essential elements needed to convey your manuscript, for example Abstract, Keywords, Introduction, Materials and Methods, Results, Conclusions, Artwork and Tables.

If your article includes any Videos and/or other Supplementary material, this should be included in your initial submission for peer review purposes.

Divide the article into clearly defined sections.

#### *Figures and tables embedded in text*

Please ensure the figures and the tables included in the single file are placed next to the relevant text in the manuscript, rather than at the bottom or the top of the file. The corresponding caption should be placed directly below the figure or table.

### **Peer review**

This journal operates a single blind review process. All contributions will be initially assessed by the editor for suitability for the journal. Papers deemed suitable are then typically sent to a minimum of two independent expert reviewers to assess the scientific quality of the paper. The Editor is responsible for the final decision regarding acceptance or rejection of articles. The Editor's decision is final. More information on types of peer review.

### **REVISED SUBMISSIONS**

#### *Use of word processing software*

Regardless of the file format of the original submission, at revision you must provide us with an editable file of the entire article. Keep the layout of the text as simple as possible. Most formatting codes will be removed and replaced on processing the article. The electronic text should be prepared in a way very similar to that of conventional manuscripts (see also the Guide to Publishing with Elsevier). See also the section on Electronic artwork.

To avoid unnecessary errors you are strongly advised to use the 'spell-check' and 'grammar-check' functions of your word processor.

### **Article structure**

#### *Subdivision - numbered sections*

Divide your article into clearly defined and numbered sections. Subsections should be numbered 1.1 (then 1.1.1, 1.1.2, ...), 1.2, etc. (the abstract is not included in section numbering). Use this numbering also for internal cross-referencing: do not just refer to 'the text'. Any subsection may be given a brief heading. Each heading should appear on its own separate line.

#### *Introduction*

State the objectives of the work and provide an adequate background, avoiding a detailed literature survey or a summary of the results.

#### *Material and methods*

Provide sufficient details to allow the work to be reproduced by an independent researcher. Methods that are already published should be summarized, and indicated by a reference. If quoting directly from a previously published method, use quotation marks and also cite the source. Any modifications to existing methods should also be described.

#### *Theory/calculation*

A Theory section should extend, not repeat, the background to the article already dealt with in the Introduction and lay the foundation for further work. In contrast, a Calculation section represents a practical development from a theoretical basis.

#### *Results*

Results should be clear and concise.

#### *Discussion*

This should explore the significance of the results of the work, not repeat them. A combined Results and Discussion section is often appropriate. Avoid extensive citations and discussion of published literature.

#### *Conclusions*

The main conclusions of the study may be presented in a short Conclusions section, which may stand alone or form a subsection of a Discussion or Results and Discussion section.

#### *Appendices*

If there is more than one appendix, they should be identified as A, B, etc. Formulae and equations in appendices should be given separate numbering: Eq. (A.1), Eq. (A.2), etc.; in a subsequent appendix, Eq. (B.1) and so on. Similarly for tables and figures: Table A.1; Fig. A.1, etc.

### **Essential title page information**

- **Title.** Concise and informative. Titles are often used in information-retrieval systems. Avoid abbreviations and formulae where possible.
- **Author names and affiliations.** Please clearly indicate the given name(s) and family name(s) of each author and check that all names are accurately spelled. You can add your name between parentheses in your own script behind the English transliteration. Present the authors' affiliation addresses (where the actual work was done) below the names. Indicate all affiliations with a lower- case superscript letter immediately after the author's name and in front of the appropriate address. Provide the full postal address of each affiliation, including the country name and, if available, the e-mail address of each author.
- **Corresponding author.** Clearly indicate who will handle correspondence at all stages of refereeing and publication, also post-publication. This responsibility includes answering any future queries about Methodology and Materials. Ensure that the e-mail address is given and that contact details are kept up to date by the corresponding author.
- **Present/permanent address.** If an author has moved since the work described in the article was done, or was visiting at the time, a 'Present address' (or 'Permanent address') may be indicated as a footnote to that author's name. The address at which the author actually did the work must be retained as the main, affiliation address. Superscript Arabic numerals are used for such footnotes.

### **Abstract**

A concise and factual abstract is required. The abstract should state briefly the purpose of the research, the principal results and major conclusions. An abstract is often presented separately from the article, so it must be able to stand alone. For this reason, References should be avoided, but if essential, then cite the author(s) and year(s). Also, non-standard or uncommon abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself.

The abstract should not exceed 200 words.

#### *Graphical abstract*

Although a graphical abstract is optional, its use is encouraged as it draws more attention to the online article. The graphical abstract should summarize the contents of the article in a concise, pictorial form designed to capture the attention of a wide readership. Graphical abstracts should be submitted as a separate file in the online submission system. Image size: Please provide an image with a minimum of  $531 \times 1328$  pixels (h  $\times$  w) or proportionally more. The image should be readable at a size of  $5 \times 13$  cm using a regular screen resolution of 96 dpi. Preferred file types: TIFF, EPS, PDF or MS Office files. You can view Example Graphical Abstracts on our information site.

Authors can make use of Elsevier's Illustration Services to ensure the best presentation of their images and in accordance with all technical requirements.

### *Highlights*

Highlights are mandatory for this journal. They consist of a short collection of bullet points that convey the core findings of the article and should be submitted in a separate editable file in the online submission system. Please use 'Highlights' in the file name and include 3 to 5 bullet points (maximum 85 characters, including spaces, per bullet point). You can view example Highlights on our information site.

### **Keywords**

Immediately after the abstract, provide a maximum of 6 keywords, using American spelling and avoiding general and plural terms and multiple concepts (avoid, for example, 'and', 'of'). Be sparing with abbreviations: only abbreviations firmly established in the field may be eligible. These keywords will be used for indexing purposes.

### **Abbreviations**

Define abbreviations that are not standard in this field in a footnote to be placed on the first page of the article. Such abbreviations that are unavoidable in the abstract must be defined at their first mention there, as well as in the footnote. Ensure consistency of abbreviations throughout the article.

### **Acknowledgements**

Collate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here those individuals who provided help during the research (e.g., providing language help, writing assistance or proof reading the article, etc.).

### *Formatting of funding sources*

List funding sources in this standard way to facilitate compliance to funder's requirements:

Funding: This work was supported by the National Institutes of Health [grant numbers xxxx, yyyy]; the Bill & Melinda Gates Foundation, Seattle, WA [grant number zzzz]; and the United States Institutes of Peace [grant number aaaa].

It is not necessary to include detailed descriptions on the program or type of grants and awards. When funding is from a block grant or other resources available to a university, college, or other research institution, submit the name of the institute or organization that provided the funding.

If no funding has been provided for the research, please include the following sentence:

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

.

### **Artwork**

- Make sure you use uniform lettering and sizing of your original artwork.
- Preferred fonts: Arial (or Helvetica), Times New Roman (or Times), Symbol, Courier.
- Number the illustrations according to their sequence in the text.
- Use a logical naming convention for your artwork files.
- Indicate per figure if it is a single, 1.5 or 2-column fitting image.
- For Word submissions only, you may still provide figures and their captions, and tables within a single file at the revision stage.
- Please note that individual figure files larger than 10 MB must be provided in separate source files.

## Formats

Regardless of the application used, when your electronic artwork is finalized, please 'save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):

EPS (or PDF): Vector drawings. Embed the font or save the text as 'graphics'.

TIFF (or JPG): Color or grayscale photographs (halftones): always use a minimum of 300 dpi.

TIFF (or JPG): Bitmapped line drawings: use a minimum of 1000 dpi.

TIFF (or JPG): Combinations bitmapped line/half-tone (color or grayscale): a minimum of 500 dpi is required.

## Figure captions

Ensure that each illustration has a caption. A caption should comprise a brief title (**not** on the figure itself) and a description of the illustration. Keep text in the illustrations themselves to a minimum but explain all symbols and abbreviations used.

Please submit tables as editable text and not as images. Tables should be placed next to the relevant text in the article. Number tables consecutively in accordance with their appearance in the text and place any table notes below the table body. Be sparing in the use of tables and ensure that the data presented in them do not duplicate results described elsewhere in the article. Please avoid using vertical rules and shading in table cells.

## References

### *Citation in text*

Please ensure that every reference cited in the text is also present in the reference list (and vice versa). Any references cited in the abstract must be given in full. Unpublished results and personal communications are not recommended in the reference list, but may be mentioned in the text. If these references are included in the reference list they should follow the standard reference style of the journal and should include a substitution of the publication date with either 'Unpublished results' or 'Personal communication'. Citation of a reference as 'in press' implies that the item has been accepted for publication and a copy of the title page of the relevant article must be submitted.

### *Reference links*

Increased discoverability of research and high quality peer review are ensured by online links to the sources cited. In order to allow us to create links to abstracting and indexing services, such as Scopus, CrossRef and PubMed, please ensure that data provided in the references are correct. Please note that incorrect surnames, journal/book titles, publication year and pagination may prevent link creation. When copying references, please be careful as they may already contain errors. Use of the DOI is highly encouraged.

A DOI is guaranteed never to change, so you can use it as a permanent link to any electronic article. An example of a citation using DOI for an article not yet in an issue is: VanDecar J.C., Russo R.M., James D.E., Ambeh W.B., Franke M. (2003). Aseismic continuation of the Lesser Antilles slab beneath northeastern Venezuela. *Journal of Geophysical Research*, <https://doi.org/10.1029/2001JB000884>. Please note the format of such citations should be in the same style as all other references in the paper.

#### *Web references*

As a minimum, the full URL should be given and the date when the reference was last accessed. Any further information, if known (DOI, author names, dates, reference to a source publication, etc.), should also be given. Web references can be listed separately (e.g., after the reference list) under a different heading if desired, or can be included in the reference list.

#### *Data references*

This journal encourages you to cite underlying or relevant datasets in your manuscript by citing them in your text and including a data reference in your Reference List. Data references should include the following elements: author name(s), dataset title, data repository, version (where available), year, and global persistent identifier. Add [dataset] immediately before the reference so we can properly identify it as a data reference. The [dataset] identifier will not appear in your published article.

#### *References in a special issue*

Please ensure that the words 'this issue' are added to any references in the list (and any citations in the text) to other articles in the same Special Issue.

#### *Reference management software*

Most Elsevier journals have their reference template available in many of the most popular reference management software products. These include all products that support Citation Style Language styles, such as Mendeley and Zotero, as well as EndNote. Using the word processor plug-ins from these products, authors only need to select the appropriate journal template when preparing their article, after which citations and bibliographies will be automatically formatted in the journal's style. If no template is yet available for this journal, please follow the format of the sample references and citations as shown in this Guide. If you use reference management software, please ensure that you remove all field codes before submitting the electronic manuscript.

When preparing your manuscript, you will then be able to select this style using the Mendeley plugins for Microsoft Word or LibreOffice.

### **Reference formatting**

There are no strict requirements on reference formatting at submission. References can be in any style or format as long as the style is consistent. Where applicable, author(s) name(s), journal title/ book title, chapter title/article title, year of publication, volume number/book chapter and the article number or pagination must be present. Use of DOI is highly encouraged. The reference style used by the journal will be applied to the accepted article by Elsevier at the proof stage.

*Text:* Indicate references by number(s) in square brackets in line with the text. The actual authors can be referred to, but the reference number(s) must always be given. Example: '..... as demonstrated [3,6]. Barnaby and Jones [8] obtained a different result ....'

*List:* Number the references (numbers in square brackets).

#### *Examples:*

##### *Reference:*

[1] J. van der Geer, J.A.J. Hanraads, R.A. Lupton, The art of writing a scientific article, *J. Sci. Commun.* 163 (2010) 51–59. <https://doi.org/10.1016/j.Sc.2010.00372>.

Reference to a journal publication with an article number:

[2] Van der Geer, J., Hanraads, J.A.J., Lupton, R.A., 2018. The art of writing a scientific article. *Heliyon*. 19, e00205. <https://doi.org/10.1016/j.heliyon.2018.e00205>.

[3] W. Strunk Jr., E.B. White, *The Elements of Style*, fourth ed., Longman, New York, 2000.

Reference to a chapter in an edited book:

[4] G.R. Mettam, L.B. Adams, How to prepare an electronic version of your article, in: B.S. Jones, R.Z. Smith (Eds.), *Introduction to the Electronic Age*, E-Publishing Inc., New York, 2009, pp. 281–304. Reference to a website:

[5] Cancer Research UK, Cancer statistics reports for the UK.

[6] M. Oguro, S. Imahiro, S. Saito, T. Nakashizuka, Mortality data for Japanese oak wilt disease and surrounding forest compositions, Mendeley Data, v1, 2015.

<https://doi.org/10.17632/xwj98nb39r.1>.

### **Journal abbreviations source**

Journal names should be abbreviated according to the List of Title Word Abbreviations.

### **Video**

Elsevier accepts video material and animation sequences to support and enhance your scientific research. Authors who have video or animation files that they wish to submit with their article are strongly encouraged to include links to these within the body of the article. This can be done in the same way as a figure or table by referring to the video or animation content and noting in the body text where it should be placed. All submitted files should be properly labeled so that they directly relate to the video file's content. . In order to ensure that your video or animation material is directly usable, please provide the file in one of our recommended file formats with a preferred maximum size of 150 MB per file, 1 GB in total. Video and animation files supplied will be published online in the electronic version of your article in Elsevier Web products, including ScienceDirect. Please supply 'stills' with your files: you can choose any frame from the video or animation or make a separate image. These will be used instead of standard icons and will personalize the link to your video data. For more detailed instructions please visit our video instruction pages. Note: since video and animation cannot be embedded in the print version of the journal, please provide text for both the electronic and the print version for the portions of the article that refer to this content.

### **Data visualization**

Include interactive data visualizations in your publication and let your readers interact and engage more closely with your research. Follow the instructions here to find out about available data visualization options and how to include them with your article.

### **Supplementary material**

Supplementary material such as applications, images and sound clips, can be published with your article to enhance it. Submitted supplementary items are published exactly as they are received (Excel or PowerPoint files will appear as such online). Please submit your material together with the article and supply a concise, descriptive caption for each supplementary file. If you wish to make changes to supplementary material during any stage of the process, please make sure to provide an updated file. Do not annotate any corrections on a previous version. Please switch off the 'Track Changes' option in Microsoft Office files as these will appear in the published version.

## Research data

This journal encourages and enables you to share data that supports your research publication where appropriate, and enables you to interlink the data with your published articles. Research data refers to the results of observations or experimentation that validate research findings. To facilitate reproducibility and data reuse, this journal also encourages you to share your software, code, models, algorithms, protocols, methods and other useful materials related to the project.

Below are a number of ways in which you can associate data with your article or make a statement about the availability of your data when submitting your manuscript. If you are sharing data in one of these ways, you are encouraged to cite the data in your manuscript and reference list. Please refer to the "References" section for more information about data citation. For more information on depositing, sharing and using research data and other relevant research materials, visit the research data page.

### 5.3.2 Estudo de Revisão Sistemática (Artigo 2)

Journal Of Pediatric Hematology And Oncology

#### Scope

*Journal of Pediatric Hematology-Oncology* reports on major advances in the diagnosis and treatment of cancer and blood diseases in children. Each issue presents informative case studies and original research articles from leading clinicians and investigators worldwide.

#### Ethical/legal considerations

A submitted manuscript must be an original contribution not previously published (except as an abstract or a preliminary report), must not be under consideration for publication elsewhere, and, if accepted, must not be published elsewhere in similar form, in any language, without the consent of Lippincott Williams & Wilkins. Each person listed as an author is expected to have participated in the study to a significant extent. Although the editors and referees make every effort to ensure the validity of published manuscripts, the final responsibility rests with the authors, not with the Journal, its editors, or the publisher. All manuscripts must be submitted on-line through the journal's Web site <https://jpho.edmgr.com/>. See submission instructions under "On-line manuscript submission."



**Patient Anonymity and Informed Consent:** It is the author's responsibility to ensure that a patient's anonymity be carefully protected and to verify that any experimental investigation with human subjects reported in the manuscript was performed with informed consent and following all the guidelines for experimental investigation with human subjects required by the institution(s) with which all the authors are affiliated. Authors should mask patients' eyes and remove patients' names from figures. Photographs with bars placed over the eyes of patients CANNOT be used in publication, unless they obtain written consent from the patients and submit written consent with the manuscript.

### **Copyright:**

The hard copy.pdf file is no longer in use. The corresponding author will complete the authorship verification questionnaire within the submission steps, and provide each co-authors email address. The co-authors are emailed a hyperlink which they will verify their co-authorship electronically. The co-authors are not required to register for an account in Editorial Manager and no new accounts are created for them, rather, they are completing a form that is tied to the submission record.

### **Conflicts of interest**

Authors must state all possible conflicts of interest in the manuscript, including financial, consultant, institutional and other relationships that might lead to bias or a conflict of interest. If there is no conflict of interest, this should also be explicitly stated as none declared. All sources of funding should be acknowledged in the manuscript. All relevant conflicts of interest and sources of funding should be included on the title page of the manuscript with the heading "Conflicts of Interest and Source of Funding:". For example:

Conflicts of Interest and Source of Funding: A has received honoraria from Company Z. B is currently receiving a grant (#12345) from Organization Y, and is on the speaker's bureau for Organization X – the CME organizers for Company A. For the remaining authors none were declared.

**Compliance with NIH and Other Research Funding Agency Accessibility Requirements:** A number of research funding agencies now require or request authors to submit the post-print (the article after peer review and acceptance but not the final published article) to a repository that is accessible online by all without charge. As a service to our authors, LWW will identify to the National Library of Medicine (NLM) articles that require deposit and will transmit the post-print of an article based on research funded in whole or in part by the National Institutes of Health, Wellcome Trust, Howard Hughes Medical Institute, or other funding agencies to PubMed Central. The revised Copyright Transfer Agreement provides the mechanism.

Permissions: Authors must submit written permission from the copyright owner (usually the publisher) to use direct quotations, tables, or illustrations that have appeared in copyrighted form elsewhere, along with complete details about the source. Any permissions fees that might be required by the copyright owner are the responsibility of the authors requesting use of the borrowed material, not the responsibility of Lippincott Williams & Wilkins.

### **Manuscript submission**

On-Line Manuscript Submission: All manuscripts must be submitted on-line through the new Web site at <https://jpho.edmgr.com/>.

First-time users: Please click the Register button from the main top menu and enter the requested information. On successful registration, you will be sent an e-mail indicating your user name and password. Print a copy of this information for future reference. Note: If you have received an e-mail from us with an assigned user ID and password, or if you are a repeat user, do not register again. Just log in. Once you have an assigned ID and password, you do not have to re-register, even if your status changes (that is, author, reviewer, or editor).

### **Authors:**

Please click the login button from the menu at the top of the page and log in to the system as an Author. Submit your manuscript according to the author instructions. You will be able to track the progress of your manuscript through the system. If you experience difficulties using the system, please contact [JPHO@ymail.com](mailto:JPHO@ymail.com). Requests for help and other questions will be addressed in the order received.

### **Preparation of Manuscript:**

Manuscripts that do not adhere to the following instructions will be returned to the corresponding author for technical revision before undergoing peer review.

### **Title Page:**

Include on the title page (a) complete manuscript title; (b) authors' full names, highest academic degrees, and affiliations; (c) name and address for correspondence, including fax number, telephone number, and e-mail address; (d) address for reprints if different from that of corresponding author; and (e) all sources of support, including pharmaceutical and industry support, that require acknowledgment.

The title page must also include disclosure of funding received for this work from any of the following organizations: National Institutes of Health (NIH); Wellcome Trust; Howard Hughes Medical Institute (HHMI); and other(s).

### **Unstructured Abstract and Key Words:**

Limit the abstract to 200 words. It must be factual and comprehensive. Limit the use of abbreviations and acronyms, and avoid general statements (eg, “the significance of the results is discussed”). List three to five key words or phrases.

**Text:**

Organize the manuscript into four main headings: Introduction, Materials and Methods, Results, and Discussion. Define abbreviations at first mention in text and in each table and figure. If a brand name is cited, supply the manufacturer's name and address (city and state/country). All forms of support, including pharmaceutical industry support, must be acknowledged in the Acknowledgment section.

**Abbreviations:**

For a list of standard abbreviations, consult the Council of Biology Editors Style Guide (available from the Council of Science Editors, 9650 Rockville Pike, Bethesda, MD 20814) or other standard sources. Write out the full term for each abbreviation at its first use unless it is a standard unit of measure.

**Clinical and laboratory observations:**

Clinical observations may include case histories that demonstrate novel findings or associations, important clinical responses when a larger study is not needed to address a specific issue, or a unique laboratory observation linked to clinical care and/or practice. Text should contain 2500 words or fewer, with a brief abstract of 100 words or fewer. Abstracts outline background, observation(s), and conclusions. Include 4 illustrations and/or tables or fewer and 20 references or fewer.

**Medical progress:**

Review articles for this section should highlight what is particularly new and novel in a field related to pediatric hematology/oncology. Text should contain 5000 words or fewer and 100 references or fewer. Shorter reviews are encouraged and preferred. Authors considering submission should consult the Editor-in-Chief.

**Morphology corner:**

This section features photographs of especially interesting blood smears, bone marrow, or other tissue specimens that highlight an important aspect of hematology/oncology. Include an introduction of 200 words or fewer, the figure(s), a conclusion of 200 words or fewer, and 6 references or fewer.

**Radiology corner:**

This section features photographs of scans of radiographic studies, such as plain radiographs, bone scans, computed tomography scans, magnetic resonance images, or other

modalities highlighting a special feature of a topic or case. Include an introduction of 200 words or fewer, the figure(s), a conclusion of 200 words or fewer, and 6 references or fewer.

**HISTORICAL INSIGHTS:** Historical insights include concise descriptions or analyses of historical importance in the field of pediatric hematology/oncology. These may include personal descriptions of historical figures, important papers, and interesting occurrences that led to advancements in pediatric hematology/oncology. Photographs and artwork are welcome. Text should contain 2500 words or fewer and include 25 references or fewer. All material should be original or carry permission for publication.

#### **Letters to the editor:**

Letters to the editor should pertain to articles published within the Journal of Pediatric Hematology/Oncology or highlight important new clinical or laboratory insights. Text should contain 500 words or fewer.

#### **Book reviews:**

Reviews of books should relate to topics relevant to pediatric hematology/oncology, including immunology and transplantation. Text should contain 1000 words or fewer.

#### **Announcements:**

Announcements should be submitted 6 months in advance of the event date and may include scheduled meetings, symposia, postgraduate courses, and other announcements of interest to specialists in pediatric hematology/oncology.

#### **References:**

The authors are responsible for the accuracy of the references. Key the references (double-spaced) at the end of the manuscript. Cite the references in text in the order of appearance. Cite unpublished data—such as papers submitted but not yet accepted for publication and personal communications, including e-mail communications—in parentheses in the text. If there are more than three authors, name only the first three authors and then use et al. Refer to the List of Journals Indexed in Index Medicus for abbreviations of journal names, or access the list at <http://www.nlm.nih.gov/tsd/serials/lji.html>. Sample references are given below:

#### **Journal Article**

1. Ang KK, Price RE, Stephens LC, et al. The tolerance of primate spinal cord to re-irradiation. *Int J Radiat Oncol Biol Phys*. 1993;25:459–464.

#### **Book Chapter**

2. Dimery IW. Chemotherapy in head and neck cancer. In: Myerhoff WI, Rice DH, eds. Otolaryngology: head and neck surgery, 2nd ed. Philadelphia: WB Saunders, 1992:1027–1045.

#### Entire Book

3. Virchow R. Cellular Pathology. Philadelphia: JB Lippincott, 1863.

#### Software

4. Epi Info [computer program]. Version 6. Atlanta, GA: Centers for Disease Control and Prevention; 1994.

#### Online Journals

5. Friedman SA. Preeclampsia: a review of the role of prostaglandins. Obstet Gynecol [serial online]. January 1988;71:22–37. Available from: BRS Information Technologies, McLean, VA. Accessed December 15, 1990.

#### Database

6. CANCERNET-PDQ [database online]. Bethesda, MD: National Cancer Institute; 1996. Updated March 29, 1996.

#### World Wide Web

7. Gostin LO. Drug use and HIV/AIDS [JAMA HIV/AIDS Web site]. June 1, 1996. Available at: <http://www.ama-assn.org/special/hiv/ethics>. Accessed June 26, 1997.

#### URL (Uniform Resource Locator)

8. (J. M. Kramer, K. Kramer [[jmkramer@umich.edu](mailto:jmkramer@umich.edu)], e-mail, March 6, 1996).

#### Figures:

##### A) Creating Digital Artwork

- Learn about the publication requirements for Digital Artwork: <http://links.lww.com/ES/A42>
- Create, Scan and Save your artwork and compare your final figure to the Digital Artwork Guideline Checklist (below).
- Upload each figure to Editorial Manager in conjunction with your manuscript text and tables.

##### B) Digital Artwork Guideline Checklist

Here are the basics to have in place before submitting your digital artwork:

Artwork should be saved as TIFF, EPS, or MS Office (DOC, PPT, XLS) files. Highresolution PDF files are also acceptable.

Crop out any white or black space surrounding the image.

Diagrams, drawings, graphs, and other line art must be vector or saved at a resolution of at least 1200 dpi. If created in an MS Office program, send the native (DOC, PPT, XLS) file.

Photographs, radiographs and other halftone images must be saved at a resolution of at least 300 dpi.

Photographs and radiographs with text must be saved as postscript or at a resolution of at least 600 dpi.

Each figure must be saved and submitted as a separate file. Figures should not be embedded in the manuscript text file.

**Remember:**

Cite figures consecutively in your manuscript.

Number figures in the figure legend in the order in which they are discussed.

Upload figures consecutively to the Editorial Manager web site and enter figure numbers consecutively in the Description field when uploading the files.

**Figure Legends:**

Include legends for all figures. They should be brief and specific, and they should appear on a separate manuscript page after the references. Use scale markers in the image for electron micrographs, and indicate the type of stain used.

**Color Figures:**

The journal accepts for publication color figures that will enhance an article. Authors who submit color figures will receive an estimate of the cost for color reproduction. If they decide not to pay for color reproduction, they can request that the figures be converted to black and white at no charge.

**Tables:**

Create tables using the table creating and editing feature of your word processing software (e.g., Word, WordPerfect). Do not use Excel or comparable spreadsheet programs. Group all tables in a separate file. Cite tables consecutively in the text, and number them in that order. Each table should appear on a separate sheet and should include the table title, appropriate column heads, and explanatory legends (including definitions of any abbreviations used). Do not embed tables within the body of the manuscript. They should be self-explanatory and should supplement, rather than duplicate, the material in the text.

**Supplemental Digital Content (SDC):**

Authors may submit SDC via Editorial Manager to LWW journals that enhance their article's text to be considered for online posting. SDC may include standard media such as

text documents, graphs, audio, video, etc. On the Attach Files page of the submission process, please select Supplemental Audio, Video, or Data for your uploaded file as the Submission Item. If an article with SDC is accepted, our production staff will create a URL with the SDC file. The URL will be placed in the call-out within the article. SDC files are not copy-edited by LWW staff, they will be presented digitally as submitted. For a list of all available file types and detailed instructions, please visit <http://links.lww.com/A142>.

#### SDC Call-outs

Supplemental Digital Content must be cited consecutively in the text of the submitted manuscript. Citations should include the type of material submitted (Audio, Figure, Table, etc.), be clearly labeled as "Supplemental Digital Content," include the sequential list number, and provide a description of the supplemental content. All descriptive text should be included in the call-out as it will not appear elsewhere in the article.

#### Example:

We performed many tests on the degrees of flexibility in the elbow (see Video, Supplemental Digital Content 1, which demonstrates elbow flexibility) and found our results inconclusive.

#### **List of Supplemental Digital Content**

A listing of Supplemental Digital Content must be submitted at the end of the manuscript file. Include the SDC number and file type of the Supplemental Digital Content. This text will be removed by our production staff and not be published.

**Style:** In general, style should be patterned after the American Medical Association Manual of Style (9th edition). Stedman's Medical Dictionary (27th edition) and Merriam Webster's Collegiate Dictionary (10th edition) should be used as standard references. Drugs and therapeutic agents should be referred to by their accepted generic or chemical names. The name should not be abbreviated. Code numbers should be used only when a generic name is not yet available. In that case, the chemical name and a figure giving the chemical structure of the drug is required. Copyright or trade names of drugs should be capitalized and placed in parentheses after the name of the drug. Names and locations (city and state in USA; city and country outside USA) of manufacturers of drugs, supplies, or equipment cited in a manuscript are required to comply with trademark law and should be provided in parentheses. Units of measure should be expressed in the metric system, and temperatures should be expressed in degrees Celsius. Conventional units should be written as SI units as appropriate.

## **6 DISCUSSÃO GERAL**



A associação entre MO em pacientes pediátricos portadores de LLA está escassamente contemplada na literatura. De maneira geral, a fisiopatologia do acometimento ocular na LLA deve-se a três mecanismos principais: I) infiltração direta do olho e órbita pelas células neoplásicas; II) anormalidades vasculares afetando a retina ou III) acometimento neuro-oftalmológico (REDDY; MENON, 1998). Não podemos ignorar ainda, os efeitos dos quimioterápicos na visão dos pacientes. O uso de altas doses de GC no tratamento desses pacientes, por si só, já implica em um risco significativamente aumentado de desenvolver, por exemplo, HO, catarata, glaucoma cortisônico, olho seco, diplopia e cegueira (MENDONÇA et al., 2014; WHELAN et al., 2010), que pode persistir em alguns casos, cessado o seu uso.

O primeiro artigo apresenta os resultados da primeira investigação com delineamento prospectivo em um estudo de coorte prospectivo em crianças e adolescentes com LLA, em um serviço público especializado em oncologia pediátrica localizado na região nordeste do Brasil entre 1º de janeiro de 2013 e 30 de dezembro de 2017. O diferencial em relação aos estudos anteriores foi a realização de exame oftalmológico em D0, D8, D28 e D6meses. Os resultados foram apresentados de maneira descritiva e avaliando se as MO podem estar associadas a fatores de risco para recaída da doença, tipo de uso de protocolo de tratamento, gênero, idade, imunofenotipagem, óbito e infiltração líquórica.

De um total de 78 pacientes admitidos no serviço entre 2013 e 2017, 67 preencheram os critérios de inclusão, porém 12 deles foram a óbito antes do D8. Cinquenta e cinco tiveram exame ocular realizado pelo menos uma vez entre o D8 e o D6meses. Destes, foram obtidos dois grupos: pacientes com exame normal (N=37) e pacientes que apresentaram alguma manifestação ocular (N=18). O presente estudo identificou 33% de pacientes com comprometimento ocular em alguma etapa do tratamento, contrastando com alguns resultados de estudos anteriores que obtiveram 15-39%, divergência que reflete a aplicação de diferentes métodos. Adicionalmente, no presente estudo, a maior proporção de alterações oculares se deve à inclusão dos casos de HO, que foram ativamente rastreados nessa coorte, mesmo na ausência de sintomas. Tiveram MO estatisticamente significativas os pacientes de alto risco para recaída, compatível com o descrito na literatura (RUSSO et al., 2008) e os pacientes tratados com protocolo de 1999, o que pode ser explicado pela dose total de GC, que é maior que a do protocolo de 2009 (BRANDALISE et al., 2010; WATANABE, 2007).

Envolvimento ocular direto ocorreu em um paciente (2%) sob a forma de infiltração leucêmica do nervo óptico, alteração observada em 7,3% dos casos em estudo anterior (KHADKA et al., 2014). Entre as alterações oftálmicas secundárias, a retina foi o tecido mais

afetado, ocorrendo hemorragia em 7,3% dos casos, proporção consistentemente semelhante ao descrito anteriormente: 8 - 8,8% (BITIRGEN et al., 2016; REDDY; MENON, 1998).

Outras alterações secundárias foram o desenvolvimento de celulite orbitária (2%) e uveíte devido a Herpes zoster (2%). Pacientes com LLA podem adquirir infecções por germes oportunistas e reativar infecções virais durante os períodos de neutropenia, com elevado risco de morte (COGAN, 1977).

Observou-se que, durante a fase de indução da remissão, quando doses elevadas de corticosteroides são utilizadas, houve aumento significativo da PIO em dez pacientes (20%), cujos valores foram compatíveis com HO ( $PIO > 21\text{mmHg}$ ). O resultado foi superior ao apresentado em estudo anterior, que obteve 16,6% (MENDONCA et al., 2014). Houve diferença estatisticamente significativa na variação da PIO em aferições realizadas durante a fase de indução da remissão, entre D8 e D6meses ( $p=0,026$ ). Esse achado representa o padrão de variação de PIO observado no glaucoma cortisônico (KERSEY; BROADWAY, 2005).

Inicialmente, a PIO aumentou consideravelmente, mas os valores pressóricos já estavam normalizados no D28 e D6meses, seja com a utilização do tratamento antiglaucomatoso (quando os valores de PIO eram compatíveis com HO), seja com o término da utilização de GC no final do tratamento. No glaucoma induzido por GC a elevação da PIO é gradual e progressiva. Então, assim como no glaucoma de ângulo aberto, os sintomas são pouco evidentes, mesmo com risco de danos às fibras nervosas ópticas (KHADKA et al., 2014). Como em estudos anteriores (COGAN, 1977; CURTO et al., 1989; KHADKA et al., 2014; MENDONCA et al., 2014; REDDY; MENON, 1998; RUSSO, 2008), neste estudo nenhum paciente com HO isolada desenvolveu sintomas, o que dificulta o diagnóstico quando não é feito o rastreamento sistemático de todos os pacientes, independentemente da existência de sintomas.

Houve quatro pacientes com BAV definitiva (7,3%), nenhum deles com HO isolada. Somente os pacientes com outras alterações secundárias da doença como hemorragia retiniana, palidez do nervo óptico, infiltração leucêmicas e uveíte evoluíram com BAV.

Assim, existe a possibilidade de MO e HO silenciosa, com o consequente risco de dano irreversível em pacientes sem alterações oculares no início do tratamento. Os pacientes classificados com AR para recidiva da doença no momento do diagnóstico e que fizeram uso do protocolo de tratamento ALL-99 foram os que mais apresentaram MO.

Devido à alta prevalência de HO encontrada no estudo da coorte, foi feito um estudo de revisão sistemática da literatura para avaliar a relação entre HO e Glaucoma em pacientes pediátricos tratados para LLA, devido ser a MO mais prevalente. Foram encontrados quatro

artigos: dois relatos de casos (PILBEAM; SALVI; HAVANI et al., 2012; THAM et al., 2004), uma série de cinco casos (YAMASHITA et al., 2010) e um estudo descritivo prospectivo de 12 pacientes (MENDONÇA et al., 2014). Foram diagnosticadas nove crianças com HO.

O primeiro estudo foi apresentado por Tham e colaboradores em 2004 relatando um caso de HO em paciente tratado para LLA, cuja PIO variou de 16 a 17 mmHg antes do uso de GC, até 52 mmHg em OD e 47 mmHg no OE no décimo dia de tratamento. Esse paciente recebeu doses elevadas de GC sistêmico e, apesar do tratamento com medicações tópicas, só houve retorno para níveis após dois dias de suspensão desta medicação. Após novo ciclo de tratamento com GC oral (dexametasona) o paciente apresentou o mesmo comportamento da PIO, com o retorno aos níveis básicos somente com a interrupção da dexametasona (THAM et al., 2004). Esse relato de caso exemplifica a real fisiopatologia da HO induzida pelo GC, caracterizada pelo retorno a níveis de PIO normais após suspensão do GC (ZHANG; CLARK; YORIO, 2005).

Yamashita et al. (2010) relataram uma série retrospectiva de casos envolvendo crianças de até seis anos de idade portadoras de LLA. Eles foram tratados com prednisolona, 60 mg/m<sup>2</sup>/dia na fase de indução da remissão, e dexametasona 6 mg/m<sup>2</sup>/dia na fase de manutenção. Todos os pacientes tiveram a pressão ocular acima de 21 mmHg e foram tratados com medicações para diminuir níveis pressóricos oculares. No final, após suspensão do GC, nenhum dos pacientes necessitou de medicações para diminuir a PIO. Um deles apresentou atrofia óptica glaucomatosa, diminuição da espessura das fibras nervosas da retina (confirmado pelo OCT), apesar de se ter obtido o controle da PIO. Nesse estudo, os cinco pacientes foram encaminhados para um oftalmologista por causa de seus sintomas oculares (YAMASHITA et al., 2010). A forma em que foi aferida a PIO, por sopro de ar ou por tonometria de não-contato, não tem a previsibilidade da tonometria de contato, resultando em possível superestimação dos valores em crianças (BRESSON-DUMONT, 2009). Este fato pode explicar o achado de HO em todos os pacientes desse estudo. Devido a estes resultados controversos, estudos controlados com medição da PIO por tonometria de aplanção se fazem necessários a fim de garantir diagnósticos mais precisos. Todos os pacientes receberam medicação contra o glaucoma, um achado oposto aos resultados da coorte *Childhood Cancer Survivor Study* (CCSS), que refere como não estatisticamente significativa a possibilidade de glaucoma em crianças após cinco ou mais anos de tratadas para câncer (WHELAN et al., 2010). Possivelmente essa diferença ocorreu por superestimação dos valores da PIO, bem

como pelas particularidades de amostragem, uma vez que todos os pacientes apresentaram sintomas oculares.

Outro relato de caso foi apresentado por Pilbeam, Salvi e Havani em 2012, que avaliaram um menino de três anos de idade com diagnóstico de LLA e que no décimo dia de indução com dexametasona 3 mg/m<sup>2</sup>, apresentou fotofobia, irritabilidade, diminuição da acuidade visual e pupilas dilatadas. O exame ocular mostrou acuidade visual em 20/100 e PIO elevada (OD 42 mmHg, OE 40 mmHg). Apesar de o tratamento para o glaucoma, o paciente apresentou cegueira permanente em OD e perda significativa de visão em OE (PILBEAM; SALVI; HAVANI, 2012).

É importante ressaltar que nos estudos citados (PILBEAM; SALVI; HAVANI, 2012; YAMASHITA et al., 2010) não houve aferição da PIO antes do tratamento com GC, o que torna impossível avaliar o aumento real dos níveis pressóricos e qual seria o controle ideal da PIO. A PIO foi aferida antes do tratamento apenas em um artigo publicado (THAM et al., 2004).

Os resultados finais descritos na literatura são divergentes quanto ao controle da PIO. Embora os pacientes descritos por Yamashita e colaboradores (2010) tenham apresentado um bom controle de PIO, houve perda de fibras nervosas da retina em um paciente. No relato de Tham e colaboradores (2004) o controle da PIO foi insuficiente para manter-se em níveis adequados durante a terapia de indução. Embora os dados sobre o controle da PIO não tenham sido descritos no estudo, o paciente evoluiu com cegueira permanente no olho direito e perda importante da visão no olho esquerdo, caracterizando o efeito deletério do GC sobre a função visual (PILBEAM; SALVI; HAVANI, 2012).

Apenas o estudo realizado por Mendonca e colaboradores (2014) mostrou medidas de PIO e da função visual no início da indução de GC, bem como medidas sistemáticas durante e após o tratamento. Os autores observaram um aumento significativo da PIO em relação ao valor basal durante a fase de indução da remissão (primeiros 28 dias de tratamento), quando doses elevadas de GC foram utilizadas. Em dois de 12 pacientes (16,7%), os níveis de PIO poderiam ter sido altos o suficiente para causar neuropatia óptica glaucomatosa e, eventualmente, cegueira, uma vez que o risco de lesão do nervo glaucomatoso está diretamente relacionado ao nível e a duração da PIO.

Os autores sugerem que a verdadeira incidência de HO pode estar subestimada, uma vez que os pacientes podem permanecer assintomáticos. Exames oftalmológicos regulares e acompanhamento adicional podem revelar risco aumentado de glaucoma em sobreviventes de câncer infantil tratados com GC, e também é importante que a PIO seja medida no início da

corticoterapia, possibilitando avaliar o aumento real dos níveis pressóricos e a leitura ideal para indicar seu controle.

Assim, existe a possibilidade de HO silenciosa, com o consequente risco de dano irreversível em pacientes sem alterações oculares no início do tratamento. Identifica-se assim a necessidade de avaliação oftalmológica sistemática e universal em crianças e adolescentes com LLA desde o início do tratamento. Propõe-se a introdução de um protocolo que contemple exame oftalmológico sistemático, incluindo aferição da PIO imediatamente após o diagnóstico de LLA e antes da introdução de GC, prosseguindo-se em D8, D28 e D6meses.

## 7 CONCLUSÃO

Os pacientes estudados com LLA são jovens, com elevada expectativa de vida, pouca ou nenhuma comorbidade, e possuem doença oncológica com elevado potencial de cura. Apresentam uma alta incidência de MO devido ao tratamento e à própria doença, podendo ser assintomáticos e até evoluírem com BAV. Aqueles classificados com AR para recidiva da doença no momento do diagnóstico e os submetidos ao protocolo de 1999 são os mais propensos a apresentar MO e essas variáveis estão fortemente associadas. A HO é a MO de maior prevalência. Poucos estudos foram encontrados correlacionando crianças com LLA e HO, com resultados variando de HO silenciosa, sem alterações visuais, até cegueira irreversível.

Portanto, é proposto um protocolo que contemple exame oftalmológico geral, incluindo a medida da PIO, imediatamente após o diagnóstico de LLA (D0) e, posteriormente, em D8, D28 e D6meses. A necessidade do exame oftalmológico após os seis primeiros meses de tratamento deve ser avaliada a partir de então. Estudos clínicos controlados futuros poderão determinar a periodicidade e quais os exames oftalmológicos após D6meses.

### 7.1 Conclusão dos Artigos

#### 7.1.1 Estudo de Coorte (Artigo 1)

“Manifestações Oculares em pacientes pediátricos com Leucemia Linfoblástica Aguda: Uma coorte de 5 anos”.

#### Objetivo específico 2:

- Identificar precocemente possíveis MO em pacientes pediátricos com LLA.
- Correlacionar se fatores prognósticos estabelecidos para recidiva de doença oncológica (tipo de protocolo de tratamento quimioterápico utilizado, gênero e infiltração do líquido por células neoplásicas) podem prever risco a desenvolver MO (artigo 1).

#### Conclusão:

- a) A amostra estudada revela possibilidade de ocorrer MO em pacientes pediátricos portadores de LLA.
- b) O presente estudo identificou 33% de pacientes com MO em alguma etapa durante os seis primeiros meses de tratamento.
- c) Aqueles submetidos ao protocolo de 1999 são os que apresentam maior risco de desenvolver MO.
- d) A HO foi a alteração ocular mais prevalente (20%). Os níveis pressóricos oculares podem atingir níveis compatíveis para perda de fibras nervosas retiniana.
- e) O tratamento precoce das MO pode prevenir em alguns casos perdas visuais.
- f) Houve quatro pacientes com BAV definitiva (7,3%), nenhum deles com HO isolada.
- g) A introdução de um protocolo de avaliação oftalmológica incluindo aferição sistemática de PIO faz-se necessário, devido a possibilidade de alterações oculares, com o consequente risco de cegueira irreversível.

#### 7.1.2 Estudo de Revisão Sistemática (Artigo 2)

Steroid-Induced Ocular Hypertensive Response in Children and Adolescents with Acute Lymphoblastic Leukemia and Non-Hodgkin Lymphoma: Systematic Review.

#### **Objetivo específico 1:**

- Descrever a produção científica nacional e internacional em forma de Revisão Sistemática da MO com maior prevalência, a HO.

#### **Conclusão:**

- a) Poucos estudos específicos sobre o assunto descritos na literatura, limitando-se a dois relatos de casos, outra série de cinco pacientes e apenas um estudo descritivo, prospectivo e sistemático de 12 casos.
- b) Os resultados variaram de HO silenciosa, sem alterações visuais, até cegueira irreversível.
- c) A verdadeira incidência de HO pode estar subestimada devido a poucos estudos prospectivos controlados na literatura científica.
- d) Faz-se necessária a realização de estudos controlados, com medidas da PIO antes do início do GC, bem como de medidas seriadas padronizadas durante e após o tratamento.

## REFERÊNCIAS

- ADHIKARY, H. P.; SELLS, R. A.; BASU, P. K. Ocular complications of systemic steroid after renal transplantation and their association with HLA. **The British journal of ophthalmology**, v. 66, n. 5, p. 290–291, 1 May 1982.
- ALEMAYEHU, W.; SHAMEBO, M.; BEDRI, A. et al. Ocular manifestations of leukaemia in Ethiopians. **Ethiop Med J**, v. 34, p. 217–224, 1996.
- ALLEMANI, C.; WEIR, H. K.; CARREIRA, H. et al. Global surveillance of cancer survival 1995–2009: analysis of individual data for 25 676 887 patients from 279 population-based registries in 67 countries (CONCORD-2). **The Lancet**, [s.l.], v. 385, n. 9972, p. 977–1010, Mar. 2015. Elsevier BV. [http://dx.doi.org/10.1016/s0140-6736\(14\)62038-9](http://dx.doi.org/10.1016/s0140-6736(14)62038-9).
- ALLEN, R. A.; STRAATSMA, B. R. Ocular involvement in leukaemia and allied disorders. **Arch Ophthalmol**, v. 66, p. 490–508, 1961.
- APONTE, E. P.; DIEHL, N.; MOHNEY, B. G. Incidence and clinical characteristics of childhood glaucoma: a population-based study. **Archives of ophthalmology**, v. 128, n. 4, p. 478–482, 2010.
- ARMALY, M. F.; BECKER, B. Intraocular pressure response to topical corticosteroids. **Federation proceedings**, v. 24, n. 6, p. 1274–1278, 1 nov. 1965.
- BALLANTYNE, A. J.; MICHAELSON, I. C. **Textbook of the Fundus of the Eye**. Williams and Wilkins: Baltimore, 1970. p. 290–292.
- BERBIS, J.; MICHEL, G.; BARUCHEL, A. et al. Cohort Profile: The French Childhood Cancer Survivor Study For Leukaemia (LEA Cohort). **International Journal Of Epidemiology**, [s.l.], v. 44, n. 1, p.49-57, 17 Mar. 2014. Oxford University Press (OUP). <http://dx.doi.org/10.1093/ije/dyu031>.
- BITIRGEN, G. ; BELVIRANLI, S. ; CALISKAN, U. et al. Ophthalmic manifestations in recently diagnosed childhood leukemia. **Eur J Ophthalmol.**, v. 26, n. 1, p. 88-91, Jan./Feb. 2016.
- BRANDALISE, S. R. ; PINHEIRO, V. R. ; AGUIAR, S. S. et al. Benefits of the intermittent use of 6-mercaptopurine and methotrexate in maintenance treatment for low-risk acute lymphoblastic leukemia in children: randomized trial from the Brazilian Childhood Cooperative Group--protocol ALL-99. **Journal of Clinical Oncology**, v. 28, n. 11, p. 1911–1918, 10 Apr. 2010.



BRESSON-DUMONT, H. Intraocular pressure measurement in children. **J Fr Ophthalmol.**, v. 32, n. 3, p. 176-181, Mar. 2009.

BRITO, P.; SILVA, S. E.; COTTA, J. S. et al. Severe ocular hypertension secondary to systemic corticosteroid treatment in a child with nephrotic syndrome. **Clinical Ophthalmology**, [s.l.], p.1675-1679, Oct. 2012. Dove Medical Press Ltda.  
<http://dx.doi.org/10.2147/opth.s36261>.

ÇAÇA, I. ; ÜNLÜ, K. ; ARI, S. et al. Unilateral Optic Disc Edema in a Patient With Acute Lymphocytic Leukemia: A Case and Review of the Literature. **Annals Of Ophthalmology**, [s.l.], v. 37, n. 4, p.303-306, 2005. Springer Nature.  
<http://dx.doi.org/10.1385/ao:37:4:303>.

CAVDAR, A. O.; GOZDASOGLU, S.; ARCASOY, A. et al. Chloroma-like ocular manifestations in Turkish children with acute myelo-monocytic leukaemia. **Lancet**, v. 1, p. 680–682, 1971.

CERDÀ-IBÁÑEZ, M.; BAYO-CALDUCH, P.; MANFREDÀ-DOMÍNGUEZ, L. et al. Acute Vision Loss As The Only Sign Of Leukemia Relapse. **Retinal Cases & Brief Reports**, v. 12, p. 10–11, 2018.

CHARIF, C. M.; BELMEKKI, M.; HAJJI, Z. et al. Ophthalmic manifestations of acute leukemia. **J Fr Ophthalmol**, v. 25, n. 1, p. 62-66, 2002.

CHAUDHURI, T.; ROY, S.; ROY, P. Ischaemic optic neuropathy induced sudden blindness as an initial presentation of acute lymphoblastic leukemia. **Indian Journal Of Medical And Paediatric Oncology**, [s.l.], v. 34, n. 4, p. 335-335, 2013. Medknow.  
<http://dx.doi.org/10.4103/0971-5851.125266>.

CLAES, K.; KESTELYN, P. Ocular manifestations of graft versus host disease following bone marrow transplantation. **Bull Soc Belge Ophthalmol**, v. 77, p. 21–26, 2000.

CLARK, A. F.; WORDINGER, R. J. The role of steroids in outflow resistance. **Experimental Eye Research**, [s.l.], v. 88, n. 4, p.752-759, Apr. 2009. Elsevier BV. <http://dx.doi.org/10.1016/j.exer.2008.10.004>

COGAN, D. G. Immunosuppression and eye disease. **Am J Ophthalmol**, v. 83, p. 777–788, 1977.

COLOMBINI, A.; BARZAGHI, A.; CASTAGNETO, M. et al. Retro-orbital late relapse in a child with leukaemia after allogeneic bone marrow transplantation. **Acta Haematol**, v. 94, p. 44–47, 1995.

CULLIS, C. M.; HINES, D. R.; BULLOCK, J. D. Anterior segment ischaemia: classification and description in chronic myeloid leukaemia. **Ann Ophthalmol**, v. 11, p. 1739–1744, 1979.

CURTO, M. L. O.; ZINGONE, A.; AQUAVIVA, A. et al. Leukaemic infiltration of the eye: results of therapy in a retrospective multicentric study. **Med Pediatr Oncol**, v. 17, p. 134–139, 1989.

DANIAS, J.; GEROMETTA, R.; GE, Y. et al. Gene Expression Changes in Steroid-Induced IOP Elevation in Bovine Trabecular Meshwork. **Investigative Ophthalmology & Visual**

**Science**, [s.l.], v. 52, n. 12, p.8636-8645, 7 nov. 2011. Association for Research in Vision and Ophthalmology (ARVO). <http://dx.doi.org/10.1167/iovs.11-7563>.

DAWSON, D. M.; ROSENTHAL, D. S.; MOLONEY, W. C. Neurologic complications of acute leukaemia in adults: changing rates. **Ann Int Med**, v. 79, p. 541–544, 1973.

DE CAMARGO, B.; DE OLIVEIRA SANTOS, M.; REBELO, M. S. et al. Cancer incidence among children and adolescents in Brazil: first report of 14 population-based cancer registries. **Int J Cancer**, v. 126, p. 715–720, 2010.

DUKE-ELDER, S. System of ophthalmology. **Cornea and conjunctiva**. CV Mosby: St Louis, 1965. p. 1190–1195. Vol V111.

ESSIG, S.; LI, Q.; CHEN, Y. et al. Risk of late effects of treatment in children newly diagnosed with standard-risk acute lymphoblastic leukaemia: a report from the Childhood Cancer Survivor Study cohort. **The Lancet Oncology**, [s.l.], v. 15, n. 8, p. 841-851, July 2014. Elsevier BV. [http://dx.doi.org/10.1016/s1470-2045\(14\)70265-7](http://dx.doi.org/10.1016/s1470-2045(14)70265-7).

GIAGOUNIDIS, A. A.; MECKENSTOCK, G.; FLACKE, S. et al. Ocular changes in renal transplant patients. **American journal of ophthalmology**, v. 63, n. 2, p. 283–289, 1 fev. 1967.

GOLDMANN, H.; SCHMIDT, T. Über Applanations tonometrie. **Oph- thalmologica**, v. 134, p. 221-242, 1857.

GORDON, K. B. ; RUGO, H. S. ; DUNCAN, J. L. et al. Ocular manifestations of leukemia: leukemic infiltration versus infectious process. **Ophthalmology**, [s.l.], p. 2293-2300, Dec. 2001.

HONG, S. S.; PARK, H. R. ; KIM, K. S. et al. Late Effects, Social Adjustment, and Quality of Life in Adolescent Survivors of Childhood Leukemia. **Journal Of Korean Academy Of Nursing**, [s.l.], v. 44, n. 1, p.55-63, 2014. Korean Society of Nursing Science (KAMJE). <http://dx.doi.org/10.4040/jkan.2014.44.1.55>.

HOVLAN, K. R.; ELLIS, P. P. Ocular changes in renal transplant patients. **Am J Ophtalmol**, v. 63, n. 2, p. 283-289.

HOWARD, S. C.; METZGER, M. L.; WILIMAS, J. A. et al. Childhood Cancer Epidemiology in Low-Income Countries. **Cancer**, v. 112, n. 3, p. 461-472, 2008.

HYMAN, C. B.; BOGLE, J. M.; BRUBAKER, C.A. et al. Central nervous system involvement by leukaemia in children. **Blood**, v. 25, p. 1–12, 1965.

KASMANN, B.; RUPRECHT, K.W. Ophthalmologic findings in graft versus host disease (GvHD). **Klin Monatsbl Augenheilkd**, v. 202, p. 491–499, 1993.

KERSEY, J. P.; BROADWAY, D. C. Corticosteroid-induced glaucoma: a review of the literature. **Eye**, [s.l.], v. 20, n. 4, p.407-416, 6 May 2005. Springer Nature. <http://dx.doi.org/10.1038/sj.eye.6701895>.

KHADKA, D.; SHARMA, A. K.; SHRESTHA, J. K. et al. Ocular manifestations of childhood acute leukemia in a tertiary level eye centre of Kathmandu, Nepal. **Nepal J Ophthalmol.**, v. 6, n. 2, p. 197-204, Jul./Dec. 2014.

KINCAID, M. C.; GREEN, W. R. Ocular and orbital involvement in leukemia. **Survey Of Ophthalmology**, [s.l.], v. 27, n. 4, p.211-232, Jan. 1983. Elsevier BV. [http://dx.doi.org/10.1016/0039-6257\(83\)90123-6](http://dx.doi.org/10.1016/0039-6257(83)90123-6).

KUWABARA, T.; AIELLO, L. Leukaemic military nodule in retina. **Arch Ophthalmol**, v. 72, p. 494–497, 1964.

JABS, D. A.; HIRST, L. W.; GREEN, W. R. et al. The eye in bone marrow transplantation. II. **Histopathology. Arch Ophthalmol**, v. 101, p. 585–590, 1983.

JAKOBIEC, F. A.; JONES, I. S. Lymphomatous plasmacytic, histiocytic, haematopoietic tumours. In: JONES, I. S.; JAKOBIEC, F. A. (eds). **Diseases of the Orbit**. Hagerstown Md, Harper and Row, 1979. p. 309 (see pp 345-348).

LEE, P.-F. The Influence of Systemic Steroid Therapy on the Intraocular Pressure\*. **American Journal Of Ophthalmology**, [s.l.], v. 46, n. 3, p.328-331, set. 1958. Elsevier BV. [http://dx.doi.org/10.1016/0002-9394\(58\)90256-3](http://dx.doi.org/10.1016/0002-9394(58)90256-3)

LIN, H. F.; DAI, M. S.; KAO, W. Y. et al. Unilateral Optic Nerve Leukaemic Infiltration with Sudden Vision Loss heralding a Systemic Relapse of Acute Lymphoblastic Leukemia. **J Med Sci**, v. 25, n. 2, p. 97-100, 2005.

LINET, M. S.; BROWN, L. M.; MBULAITEYE, S. M. et al. International long-term trends and recent patterns in the incidence of leukemias and lymphomas among children and adolescents ages 0–19 years. **Int J Cancer**, v. 138, p. 1872–1874, 2015.

LINS, M. M. ; SANTOS, M. O. ; DE ALBUQUERQUE, M. F. P. M. et al. Incidence and survival of childhood leukemia in Recife, Brazil: A population-based analysis. **Pediatric Blood & Cancer**, [s.l.], v. 64, n. 8, p.1-6, 21 Dec. 2016. Wiley-Blackwell. <http://dx.doi.org/10.1002/pbc.26391>.

MARCOUX, S.; DROUIN, S. ; LAVERDIÈRE, C. et al. The PETALE study: Late adverse effects and biomarkers in childhood acute lymphoblastic leukemia survivors. **Pediatric Blood & Cancer**, [s.l.], v. 64, n. 6, p.1-8, 4 Dec. 2016. Wiley-Blackwell. <http://dx.doi.org/10.1002/pbc.26361>.

MARGOLIN, J.; RABIN, K. R.; STEUBER, C. P. et al. Acute lymphoblastic leukemia. In: PIZZO, P.A.; POPLACK, D. G. eds. **Principles and Practice of Pediatric Oncology**. 6<sup>th</sup> ed. Philadelphia, PA: Wolters Kluwer Health/Lippincott Williams & Wilkins, 2010. p. 518-565.

MCLEAN, J. M. Use of ACTH and cortisone. **Trans Am Ophthalmol Soc**, v. 48, p. 293-296, 1950.

MENDONÇA, C. Q. ; DE SOUZA, C. P.; MARTINS-FILHO, P. R. et al. Steroid-induced ocular hypertensive response in children and adolescents with acute lymphoblastic leukemia and non-hodgkin lymphoma. **Pediatric Blood & Cancer**, [s.l.], v. 61, n. 11, p.2083-2085, 17 Aug. 2014. Wiley-Blackwell. <http://dx.doi.org/10.1002/pbc.25070>

MEREDITH, T. A.; AABERG, T. M.; REESER, F. H. Rhegmatogenous retinal detachment complicating CMV retinitis. **Am J Ophthalmol**, v. 87, p. 793–796, 1979.

MERTENS, A. C.; BRAND, S.; NESS, K. et al. Health and well-being in adolescent survivors of early childhood cancer: a report from the Childhood Cancer Survivor Study. **Psycho-oncology**, [s.l.], v. 23, n. 3, p.266-275, 4 out. 2013. Wiley-Blackwell. <http://dx.doi.org/10.1002/pon.3414>.

MORENO, F.; LORIA, D.; ABRIATA, G. et al. Childhood cancer: incidence and early deaths in Argentina, 2000–2008. **Eur J Cancer**, v. 49, p. 465–473, 2013.

NG, P. C.; LEE, C. H. ; TAM, B. S. et al. Transient Increase in Intraocular Pressure during a Dose-Tapering Regime of Systemic Dexamethasone in Preterm Infants. **Ophthalmology**, [s.l.], v. 115, n. 5, p.7-14, maio 2008. Elsevier BV. <http://dx.doi.org/10.1016/j.ophtha.2008.01.010>

NGUYEN, H. S.; HAIDER, K. M.; ACKERMAN, L. L. Unusual causes of papilledema: Two illustrative cases. **Surg Neurol Int**, v. 4, p. 60, 2013.

PERRY, H. D.; MALLIN, F. J. Iris involvement in granulocytic sarcoma. **Am J Ophthalmol** v. 87, p. 530–532, 1979.

PFEFFER, B. A.; DeWITT, C. A. ; SALVADOR-SILVA, M. et al. Reduced Myocilin Expression in Cultured Monkey Trabecular Meshwork Cells Induced by a Selective Glucocorticoid Receptor Agonist: Comparison with Steroids. **Investigative Ophthalmology & Visual Science**, [s.l.], v. 51, n. 1, p.437-446, 1 jan. 2010. Association for Research in Vision and Ophthalmology (ARVO). <http://dx.doi.org/10.1167/iovs.09-4202>.

PILBEAM, K.; SALVI, S.; HAVANI, A. Corticosteroid-induce glaucoma as a complication of induction therapy in a child with acute lymphoblastic leukemia (ALL). **Pediatr Blood Cancer**, v. 58, p. 1014-1097, 2012.

POMORSKA, M.; KRZYŻANOWSKA-BERKOWSKA, P.; MISIUK-HOJŁO, M. et al. Application of optical coherence tomography in glaucoma suspect eyes. **Clinical And Experimental Optometry**, [s.l.], v. 95, n. 1, p.78-88, 10 Oct. 2011. Wiley-Blackwell. <http://dx.doi.org/10.1111/j.1444-0938.2011.00654.x>.

QASHTON, A. P. Advances in Ocular Hypertension Research and Treatment: 2011 Edition, **Scholarly Brief**, Atlanta, Georgia: Edition, ScholarlyBrief, 2011. p. 68-69.

RAVINDRANATHAN, M. P.; PAUL, V. J.; KURIAKOSE, E. T. Cataract after busulphan treatment. **Br Med J**, v. 1, n. 218–219, 1972.

RAZEGHINEJAD, M. R.; KATZ, L. J. Steroid-Induced Iatrogenic Glaucoma. **Ophthalmic Research**, [s.l.], v. 47, n. 2, p. 66-80, 2012. <http://dx.doi.org/10.1159/000328630>

REESE, A. B.; GUY, L. Exophthalmos in leukaemia. **Am J Ophthalmol**, v.16, p. 718–720, 1933.

REDDY, S. C.; MENON, B. S. A prospective study of ocular manifestations in childhood acute leukaemia. **Acta Ophthalmologica Scandinavica**, [s.l.], v. 76, n. 6, p.700-703, Dec. 1998. Wiley-Blackwell. <http://dx.doi.org/10.1034/j.1600-0420.1998.760614.x>

RIDGWAY, E. W.; JAFFE, N.; WALTON, D. S. Leukemic ophthalmopathy in children. **Cancer**, v. 38, p. 1744–1746, 1976.

ROWAN, P. J.; SLOAN, J. B. Iris and Anterior chamber involvement in leukaemia. **Ann Ophthalmol**, v. 8, p. 1081–1085, 1976.

RUSSO, A.; RIVA, I.; PIZZOLANTE, T. et al. Latanoprost ophthalmic solution in the treatment of open angle glaucoma or raised intraocular pressure: a review. **Clin Ophthalmol**, v. 2, n. 4, p. 897–905, Dec. 2008.

SCHWARTZ, B.; ROM, M. E. ; BEALKA, N. et al. Dose Response of Dexamethasone for the Enhanced Ocular Hypotensive Response to Epinephrine in Rabbits With Prior Dexamethasone Treatment. **Journal Of Ocular Pharmacology And Therapeutics**, [s.l.], v. 18, n. 2, p.133–139, Apr. 2002. Mary Ann Liebert Inc.  
<http://dx.doi.org/10.1089/108076802317373897>

SHARMA, T.; GREWAL, J. ; GUPTA, S. et al. Ophthalmic manifestations of acute leukaemias: the ophthalmologist's role. **Eye**, [s.l.], v. 18, n. 7, p. 663–672, 27 Feb. 2004. Springer Nature. <http://dx.doi.org/10.1038/sj.eye.6701308>.

SHAW, R. K.; MOORE, E. W.; FREIREICH, E. J. et al. Meningeal leukaemia. **Neurology**, v. 10, p. 823–833, 1960.

SHIBATA, K.; SHIMAMOTO, Y.; NISHIMURA, T. et al. Ocular manifestation in adult T cell leukaemia/lymphoma. **Ann Hematol**, v. 74, p. 163–168, 1997.

SHURIN, S. B.; REKATE, H. L.; ANNABLE, W. Optic atrophy induced by vincristine. **Pediatrics**, v. 70, p. 288–291, 1982.

SHUYUAN, L. Y. U.; MING, Z.; YUNXIA, G. A. O. Acute bilateral retina hemorrhages beneath internal limiting membrane: An unusual ophthalmological case report of acute leukemia during complete clinical remission. **Medicine**, v. 97, n. 7, p.e0000, Feb. 2018.

SOYLU, M.; OZDEMIR, G.; ANLI, A. Pediatric uveitis in southern Turkey. **Ocul Immunol Inflamm**, v. 5, p. 197–202, 1997.

SPAETH, G. L.; BARROS, D. S. M.; FUDEMBERG, S. J. Visual Loss Caused by Corticosteroid-Induced Glaucoma: How to Avoid It. **Retina**, [s.l.], v. 29, n. 8, p.1057–1061, Sept. 2009. Ovid Technologies (Wolters Kluwer Health).  
<http://dx.doi.org/10.1097/iae.0b013e3181b32cfd>.

STAMER, W. D. ; HOFFMAN, E. A. ; KURALI, E. et al. Unique Response Profile of Trabecular Meshwork Cells to the Novel Selective Glucocorticoid Receptor Agonist, GW870086X. **Investigative Ophthalmology & Visual Science**, [s.l.], v. 54, n. 3, p. 2100–2107, 27 Mar. 2013. Association for Research in Vision and Ophthalmology (ARVO).  
<http://dx.doi.org/10.1167/iovs.12-11298>

TALCOTT, K. E.; GARG, R. J.; GARG, S. J. Ophthalmic manifestations of leukemia. **Current Opinion In Ophthalmology**, [s.l.], v. 27, n. 6, p.545–551, Nov. 2016. Ovid Technologies (Wolters Kluwer Health).  
<http://dx.doi.org/10.1097/icu.0000000000000309>

TERWILLIGER, T.; ABDUL-HAY, M. Acute lymphoblastic leukemia: a comprehensive review and 2017 update. **Blood Cancer Journal**, [s.l.], v. 7, n. 6, p.577-577, 30 Jun. 2017. Springer Nature. <http://dx.doi.org/10.1038/bcj.2017.53>.

THAM, C. C. Y. ; NG, J. S. K. ; LI, R. T. H. et al. Intraocular pressure profile of a child on a systemic corticosteroid. **American Journal Of Ophthalmology**, [s.l.], v. 137, n. 1, p.198-201, Jan. 2004. Elsevier BV. [http://dx.doi.org/10.1016/s0002-9394\(03\)00838-9](http://dx.doi.org/10.1016/s0002-9394(03)00838-9).

TRIPATHI, R. C. ; KIRSCHNER, B. S. ; KIPP, M. et al. Corticosteroid treatment for inflammatory bowel disease in pediatric patients increases intraocular pressure. **Gastroenterology**, v. 102, n. 6, p. 1957–1961, 1992.

URBAN, R. C.; DREYER, E. B. Corticosteroid-induced glaucoma. **International Ophthalmology Clinics**, v. 33, n. 2, p. 135–139, 1993.

WALTON, R. C.; REED, K. L. Herpes zoster ophthalmicus following bone marrow transplantation in children. **Bone Marrow Transplant**, v. 23, p. 1317–1320, 1999.

WARD, E.; DESANTINS, C. ; ROBBINS, A. et al. Childhood and adolescent cancer statistics, 2014. **Ca: A Cancer Journal for Clinicians**, [s.l.], v. 64, n. 2, p.83-103, 31 Jan. 2014. American Cancer Society. <http://dx.doi.org/10.3322/caac.21219>.

WATANABE, A. Recent advance in treatment of childhood acute lymphoblastic leukemia. **Gan to Kagaku Ryoho**, v. 34, p.150-155, 2007.

WHELAN, K. F.; STRATTON, K.; KAWASHIMA, T. et al. Ocular late effects in childhood and adolescent cancer survivors: A report from the childhood cancer survivor study. **Pediatric Blood & Cancer**, [s.l.], v. 54, n. 1, p. 103-109, Jan. 2010. Wiley-Blackwell. <http://dx.doi.org/10.1002/pbc.22277>.

YALCINBAYIR, O.; BAYTAN, B. ; GELISKEN, O. et al. Spectral domain optical coherence tomography findings of patients under treatment for pediatric acute lymphoblastic leukemia. **Journal Of American Association For Pediatric Ophthalmology And Strabismus**, [s.l.], v. 21, n. 2, p.131-135, Apr. 2017. Elsevier BV. <http://dx.doi.org/10.1016/j.jaapos.2016.12.002>.

YAMASHITA, T.; KODAMA, Y.; TANAKA, M. et al. Steroid-induced Glaucoma in Children With Acute Lymphoblastic Leukemia. **Journal Of Glaucoma**, [s.l.], v. 19, n. 3, p.188-190, Mar. 2010. Ovid Technologies (Wolters Kluwer Health). <http://dx.doi.org/10.1097/ijg.0b013e3181af321d>

YASMEEN, N.; ASHRAF, S. Childhood acute lymphoblastic leukaemia; epidemiology and clinicopathological features. **JPMA: The Journal of the Pakistan Medical Association**, v. 59, n. 3, p. 150–153, 2009.

YEOH, A. J. ; TAN, D. ; LI, C. K. et al. Management of adult and paediatric acute lymphoblastic leukaemia in Asia: resource-stratified guidelines from the Asian Oncology Summit 2013. **The Lancet Oncology**, [s.l.], v. 14, n. 12, p. 508-523, Nov. 2013. Elsevier BV. [http://dx.doi.org/10.1016/s1470-2045\(13\)70452-2](http://dx.doi.org/10.1016/s1470-2045(13)70452-2).

YOUNG, J. L.; RIES, L. G.; SILVERBERG, E. et al. Cancer incidence, survival, and mortality for children younger than age 15 years. **Cancer**, v. 58, n. 2, p. 598-602, 1986.

ZHANG, X.; CLARK, A. F.; YORIO, T. Regulation of Glucocorticoid Responsiveness in Glaucomatous Trabecular Meshwork Cells by Glucocorticoid Receptor- $\beta$ . **Investigative Ophthalmology & Visual Science**, [s.l.], v. 46, n. 12, p.62-66, 1 Dec. 2005. Association for Research in Vision and Ophthalmology (ARVO). <http://dx.doi.org/10.1167/iovs.05-0571>.

### PRODUÇÃO CIENTÍFICA

- a) **“Steroid-Induced Ocular Hypertensive Response in Children and Adolescents with Acute Lymphoblastic Leukemia and Non-Hodgkin Lymphoma”.**

Artigo publicado no “Pediatric Blood and Cancer em 25/03/2014, PBC 25070

- b) **“Ocular Manifestations in Acute Lymphoblastic Leukemia: A Five-Year Cohort Study of Pediatric Patients”.**

Artigo publicado “Leukemia Research, em 29/11/2018, leukres 10.1016.

- c) **“Steroid-Induced Ocular Hypertensive Response in Children and Adolescents with Acute Lymphoblastic Leukemia and Non-Hodgkin Lymphoma: a Systematic Review”.**

Artigo submetido no “Journal of Pediatric Hematology and Oncology JPHO-18-150.

Foram submetidas as correções solicitadas pelos revisores; manuscrito com o Editor, aguardando parecer final.

- d) **“Comportamento Da Pressão Ocular Em Pacientes Pediátricos Tratados Para Leucemia Linfoblástica Aguda E Linfoma Não Hodgkin”.**

Apresentado no Congresso Brasileiro de Hematologia, Hemoterapia e Terapia Celular -Hemo 2014.

- e) **“Ocular Hypertension In Pediatric Patients Treated for Acute Lymphoblastic Leukemia and Non-Hodkin Lymphoma”**

Apresentado no 48º Congress of the International Society of Pediatric Oncology, Dublin-Irlanda, SIOP 2016.

f) **“Manifestações Oculares Em Leucemia Linfoblástica Aguda: Estudo De Coorte Em Pacientes Pediátricos Durante Cinco Anos”.**

Apresentado no Congresso Brasileiro de Hematologia, Hemoterapia e Terapia Celular -Hemo 2018.

g) **“Dissertação de Mestrado”**

- Cristiano de Queiroz Mendonça - UFS.
- Marcelle Vieira Freire - UFS.

h) **“Retinopatia Leucêmica: Relato de Caso”.**

Apresentado no IX Congresso Nacional da Sociedade Brasileira de Oftalmologia – 2017

i) **“Trabalhos de Conclusão de Curso (TCC)”:**

- Celso de Souza Dias Júnior - UFS.
- Cristiano Prado de Souza Júnior - UFS.
- Mayo Kayann Guerra Silva Tavares - UFS.
- Marcelle Vieira Freire - UFS.
- Marcus Vinícius Prado Guerreiro Filho - UNIT.
- Paulo Cesar Machado Filho - UNIT.
- Rafael Sucupira Menezes - UNIT.
- Wallace Marcelo Almeida Silva - UFS.



## **APÊNDICE A - Termo de Consentimento Livre e Esclarecido**

Prezado(a) Sr(a) \_\_\_\_\_

Meu nome é Cristiano de Queiroz Mendonça, sou Médico Oftalmologista, aluno do doutorado em Ciências da Saúde da Universidade Federal de Sergipe, tendo como orientadora no projeto de pesquisa a Professora Doutora Rosana Cipolotti.

Em virtude de poucos estudos sobre as alterações oftalmológicas e a alta prevalência dessa patologia no Brasil, fica evidente a necessidade de novas pesquisas para melhoria da qualidade de vida destes pacientes.

Assim, através deste documento, estou solicitando ao Sr (a) a liberação dos resultados dos exames que estão nos prontuários regularmente cadastrados no Ambulatório de Oncologia Pediátrica, bem como a solicitação para a realização de exame de medida da pressão intra ocular por método não invasivo de Tonometria de Aplanção, com instilação de gotas de colírio com efeito anestésico e corante de Fluoresceína. A curvatura e elevação corneana por imagem obtida pelo aparelho Pentacam HR e a análise das fibras nervosas retinianas por aparelho de Tomógrafo de Coerência Óptica, sem contato direto ocular e posterior análise em programas específicos dos aparelhos. Garantindo-se que não haverá qualquer identificação do nome do (a) paciente, afirmo desde já, que se trata de um trabalho de pesquisa e como tal, a

sua participação é voluntária, estando garantido o seu direito ao tratamento em qualquer situação. Além disso, mesmo concordando, o (a) senhor (a) pode desistir a qualquer momento, sem prejuízo para o tratamento.

Em caso de dúvida, entre em contato imediatamente conosco, nos ambulatórios onde costuma ser atendido, ou por um dos telefones anotados abaixo. Muito obrigado.

Responsável: \_\_\_\_\_

\_\_\_\_\_  
Dr Cristiano de Queiroz Mendonça Tel: 981019901

Doutorando

\_\_\_\_\_  
Drª Rosana Cipolotti Tel: 99811238

Orientadora

Data: \_\_\_\_\_

### **ANEXO A – PARECER CONSUBSTANCIADO CEP/UFS**

Hospital Universitário / Universidade Federal de Sergipe

Parecer consubstanciado do CEP

Dados do projeto de pesquisa título da pesquisa: comportamento da pressão ocular em pacientes pediátricos

Tratados para leucemia linfoblástica aguda e linfoma não hodgkin pesquisador: Cristiano MENDONÇA

Área temática: versão: 1caae: 13317113.0.0000.5546 instituição proponente: fundacao universidade federal de sergipe patrocinador principal: financiamento próprio

Dados do parecer número do parecer: 214.759

Data da relatoria: 01/03/2013

#### Apresentação do projeto:

Trata-se de estudo de pesquisa do tipo descritivo, observacional, longitudinal a ser realizado em crianças e adolescentes, ambos os sexos, na faixa etária entre 05 e 18 anos de idade, com diagnóstico de Ila e Inh, matriculados no centro de oncologia de sergipe dr. Oswaldo leite como tratamento inicial. A inclusão se dará por confirmação cito-histopatológica do diagnóstico e consentimento do responsável. A amostra será de 50 pacientes por um período de 12 meses de coleta. Apresenta critérios de inclusão e critérios de exclusão.

Os dados obtidos serão armazenados em banco de dados padronizado. Os descritivos serão expressos em medidas de tendência central (média e desvio-padrão ou mediana). As comparações entre os grupos serão feitas pelos testes de proporções (qui-quadrado ou exato de fisher) e de testes de médias (anova, teste t para amostras independentes ou Pareadas) e análise multivariada para  $p > 0,05$ .

#### Objetivo da pesquisa:

Objetivo primário: avaliar o comportamento da pressão intraocular em pacientes pediátricos portadores de neoplasias linfoproliferativas agudas tratados com glicocorticóides, nas diferentes fases do protocolo terapêutico. <sup>[11]</sup>Objetivo secundário: associar os achados com variáveis metabólicas relacionadas ao uso de glicocorticoides. Identificar eventuais interferências sobre a função visual.

Endereço: rua cláudio batista s/no bairro: sanatório

Cep: 49.060-110 e-mail: cephu@ufs.br

Uf: se município: telefone: (79) 2105-1805

Aracaju

#### Avaliação dos riscos e benefícios:

Os riscos foram considerados desprezíveis pelos pesquisadores, limitado ao desconforto da aplicação de colírio anestésico associado ao corante de fluoresceína. Considerado risco mínimo de infecção, semelhante a da instilação ocular de qualquer medicamento benefícios: diagnosticar e evitar quaisquer alterações visuais consequente a hiperetensão ocular

secundária ao tratamento para leucemia linfoblástica aguda e linfoma não hodgkin

Comentários e considerações sobre a pesquisa:

O projeto de pesquisa apresenta delineamento adequado com base na literatura científica. Método, contempla o desenho de estudo em todos os seus aspectos, local de estudo, seleção da amostra, critérios de inclusão e exclusão, procedimentos da coleta de dados e seus respectivos instrumentos e, a análise estatística.

Cronograma adequado ao estudo. Orçamento, planilha identifica os custos da pesquisa e declara ser de responsabilidade dos pesquisadores apesar de seu custo bastante expressivo (R\$ 37.705,00), assim descrito: o grupo de pesquisa já possui os equipamentos pentacam e tomógrafo de coerência óptica, sendo os exames realizados sem custo adicional. Os exames de glicemia fazem parte do protocolo de tratamento e são custeados pelo SUS. O material de consumo será custeado pelos pesquisadores.

Considerações sobre os termos de apresentação obrigatória:

Folha de rosto: preenchimento adequado pelo pesquisador e pela instituição proponente núcleo de pós-graduação em medicina. autorização do centro de oncologia de Sergipe Dr. Osvaldo Leite do Hospital de Urgências de Sergipe, datada e assinada pelo responsável

TCLE atende o que determina a Res 196/96/CNS: linguagem clara com as informações pertinentes, objetivos, riscos e benefícios, direito de desistir da participação sem prejuízo ao tratamento, o sigilo, o caráter de confiabilidade estão assegurados, contatos dos pesquisadores.

Recomendações:

Sugiro correção no TCLE identificar local para assinatura do responsável ou representante legal

Conclusões ou pendências e lista de inadequações:

TCLE identificar local para assinatura do responsável ou representante legal

Situação do parecer:

Aprovado

Necessita apreciação da conep:

Não

Endereço: rua cláudio batista s/no bairro: sanatório

Cep: 49.060-110<sup>UF</sup><sub>SE</sub>e-mail: cephu@ufs.br

Uf: se município: telefone: (79)2105-1805

Aracaju